

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
12 May 2005 (12.05.2005)

PCT

(10) International Publication Number
WO 2005/042464 A1

(51) International Patent Classification⁷: **C07C 205/11**,
255/50, 323/25, C07D 213/74, A61K 31/04, 31/435,
31/277, 31/10, A61P 5/28

(74) Agents: **BANNERMAN, David, Gardner** et al.; With-
ers & Rogers, Goldings House, 2 Hays Lane, London SE1
2HW (GB).

(21) International Application Number:
PCT/GB2004/004464

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(22) International Filing Date: 21 October 2004 (21.10.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0324551.1 21 October 2003 (21.10.2003) GB

(71) Applicant (*for all designated States except US*): **KARO
BIO AB** [SE/SE]; Novum, S-141 57 Huddinge (SE).

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for MG only*): **ELSY, David** [GB/GB]; With-
ers & Rogers, Goldings House, 2 Hays Lane, London SE1
2HW (GB).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): **JERNSTEDT,
Henrik** [SE/SE]; Egilsgatan 13A, S-753 35 Uppsala (SE).
GARG, Neeraj [IN/SE]; Barkvägen 15, S-147 52 Tumba
(SE). **GUSTAVSSON, Annika** [SE/SE]; Koltrastvägen
26, S-178 39 Ekerö (SE). **GILLNER, Mikael** [SE/SE];
Renstiernas Gata 38, 3tr, S-116 31 Stockholm (SE).
GARCIA COLLAZO, Ana, Maria [—/SE]; Möregatan
10, 6tr, S-118 27 Stockholm (SE). **KOCH, Eva** [SE/SE];
Brunbärsvägen 8, S-114 21 Stockholm (SE).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: ANDROGEN RECEPTOR MODULATORS



(I)

(57) Abstract: Treatment of Diseases caused by Disturbances of the Activity of the Androgen Receptor uses of compounds of Formula (I): (as defined herein), for the treatment of diseases caused by disturbances of the activity of androgen receptor are provided: Formula (I). Isolated compounds of Formula (I) are also provided.

WO 2005/042464 A1

NOVEL COMPOUNDS

Field of Invention

This invention relates to novel compounds which are androgen receptor ligands, to methods of preparing such compounds and to methods for using such compounds such as for androgen hormone replacement therapy and for diseases modulated by the androgen receptor such as benign prostatic hyperplasia, prostate cancer, alopecia, hirsutism, bone loss, bone fractures, osteoporosis, cachexia, and muscle wasting.

Background of Invention

The androgen receptor (AR) is a member of the steroid hormone nuclear receptor family of ligand activated transcription factors. This group includes estrogen, progesterone, mineralocorticoid, and glucocorticoid receptors all of which are activated by endogenous steroid hormones to control the expression of responsive genes. The hormone receptors share a modular structure consisting of a variable amino-terminal domain (NTD), a highly conserved DNA-binding domain (DBD), and a carboxy-terminal ligand-binding domain (LBD). The DNA-binding domain generates much of the transcriptional specificity due to its ability to discern different DNA response elements with the promoter regions of target genes. The LBD is required for ligand dependent transcriptional activity containing both the hormone-binding pocket and an important transcriptional activation functional region (AF2) required for recruitment of coactivators and the cellular transcriptional machinery.

Regulation of nuclear receptor activity resides predominantly in the binding of the

hormone ligand within the LBD. The amino acids lining the interior of the hormone-binding cavity define the selectivity of the receptor for its hormone. This allows AR to discriminate between the natural ligands and non-natural ligands.

Another level of transcriptional control is conveyed by the nuclear receptor's environment. It is widely accepted that different effector proteins (coactivators and corepressors) exist within different cell types and can lead to different patterns of gene expression. Because the conformational state of the receptor dictates which coactivator is recruited in a given cell type, it also imparts transcriptional selectivity. It is precisely this type of control that gave rise to tissue selective receptor modulators. For example, tamoxifen is a prototypical estrogen receptor selective modulator with differing properties within breast and uterine tissues. Exploitation of the conformational changes induced by synthetic ligands within the hormone-binding cavity has led to multiple generations of tissue selective receptor modulators for the estrogen receptor and can be applied to developing modulators of other nuclear receptors such as the androgen receptor.

The use of natural and synthetic androgen in hormone replacement therapy has been shown to markedly decrease the risk of osteoporosis and muscle wasting. In addition, there is evidence that hormone replacement therapy has cardiovascular benefits. However hormone replacement therapy is also associated with an increase risk of prostate cancer. It is known that certain types of synthetic AR ligands display a mixed agonist/antagonist profile of activity showing agonist activity in some tissues and antagonist activity in other tissues. Such ligands are referred to as selective androgen receptor modulators (SARMS).

What is needed in the art are compounds that can produce the same positive responses as androgen replacement therapy without the negative side effects. Also needed are androgen-like compounds that exert selective effects on different tissues of the body.

The amino acids and the “space” they define as the hormone-binding cavity can be exploited in synthesizing modulators that are highly receptor selective. These interactions between the endogenous hormone and amino acid residues within the ligand-binding cavity induce conformational changes that are distributed throughout the entire receptor structure. It is these conformational changes that lead to the dissociation of chaperone proteins that stabilize the receptors in the absence of ligand and the association of coactivator proteins. A liganded receptor devoid of its chaperone proteins is able to dimerize, translocate, recruit coactivators, and initiate transcription.

The natural ligand for the androgen receptor, androgen, is produced in both men and women by the gonads, adrenal glands and locally in target tissues. The levels of androgens secreted by the gonads are tightly regulated by a feedback mechanism involving the hypothalamus and pituitary.


In men, androgens are necessary for masculinization and fertility. However, systemic androgen excess causes testicular atrophy and infertility. Androgens may also contribute to lipid abnormalities, cardiovascular disease and psychological abnormalities. Local androgen excess is implicated in the pathogenesis of male pattern baldness (alopecia), benign prostatic hyperplasia (BPH) and acne. The physiologic role of androgens in women is not well understood, but these steroids do play a role in the development of normal body hair and libido. In women, relative androgen excess causes hirsutism (excessive hair growth), amenorrhea (abnormal loss or suppression of menses), acne and male pattern baldness.

The risk of developing prostate cancer increases dramatically with age. More than 75% of prostate cancer diagnoses are in men over the age of 65, and the prevalence of clinically undetectable prostate cancer in men over 80 years old is as high as 80%. It remains unclear as to the exact cause of prostate cancer, however, it is widely accepted that androgens can increase the severity and the rate of progression of the disease.

Androgen deprivation therapy has been the basis for prostate cancer therapy since 1941 when castration was shown to have beneficial effects on advanced stages of the disease. Hormonal intervention is currently based on disrupting the hypothalamus-pituitary-gonadal feedback mechanism to control the levels of endogenous androgens from the testes. Antiandrogens are incorporated in later stage therapies to work at the level of the androgen receptor itself, blocking residual androgens from adrenal sources. In spite of these treatments, there exists a need for an improved therapy of diseases linked to disturbances in the activity of the androgen receptor.

SUMMARY OF THE INVENTION

The present invention provides the use of a compound according to Formula I for the preparation of a medicament, wherein Formula I is defined as:



Formula I

in which;

R₁ and R₂ are the same or different and independently selected from the group consisting of; hydrogen, halogen, C₁-C₁₀ alkyl, C₁-C₁₀ substituted alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkoxy, C₁-C₁₀ alkenoxy, C₁-C₁₀ alkynoxy, C₁-C₁₀ alkylthio, C₁-C₁₀ alkenylthio, C₁-C₁₀ alkynylthio, C₆-C₁₀ arylthio, C₁-C₁₀ alkylsulphone, C₁-C₁₀ alkenylsulphone, C₁-C₁₀ alkynylsulphone, C₆-C₁₀ arylsulphone, C₁-C₁₀ alkylsulphoxide, C₁-C₁₀ alkenylsulphoxide, C₁-C₁₀ alkynylsulphoxide, C₆-C₁₀ arylsulphoxide, C₁-C₁₀ alkylarylthio, C₁-C₁₀ alkylarylulphone, C₁-C₁₀ alkylarylulphoxide, C₆-C₁₀ aryl, or C₅-C₂₀ heteroaryl, optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be the same or different; or R₁ and R₂ may together form a C₃-C₁₀ cycloalkyl group;

R₃ and R₄ are the same or different and independently selected from hydrogen, halogen, C₁-C₂₀ alkyl, C₃-C₇ cycloalkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ alkenoxy, C₁-C₄ alkynoxy, C₁-C₄ alkylthio, C₁-C₄ alkenylthio, C₁-C₄ alkynylthio, C₁-C₁₀ alkylsulphone, C₁-C₁₀ alkenylsulphone, C₁-C₁₀ alkynylsulphone, C₆-C₁₀ arylsulphone, C₁-C₁₀ alkylsulphoxide, C₁-C₁₀ alkenylsulphoxide, C₁-C₁₀ alkynylsulphoxide, C₆-C₁₀ arylsulphoxide, C₁-C₁₀ alkylarylthio, C₁-C₁₀ alkylarylulphone, C₁-C₁₀ alkylarylulphoxide, C₆-C₁₅ aryl, C₅-C₂₀ heteroaryl optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be the same or different; or can together form a keto group;

R₅ is chosen from the group consisting of; nitro, cyano, -CH₂CN, -COMe, acetic acid, halogen, sulphonic acid, -SO₂CH₃, aldehyde, carboxylic acid or ester, phosphonic acid or ester;

R₆ is chosen from the group consisting of; hydrogen, C₁-C₅ alkyl, halogen, CN, CO₂H, CHF₂, CH₂F or CF₃;

R₇ is chosen from the group consisting of; H, halogen or C₁-C₅ alkyl;

R₈ is chosen from the group consisting of; hydrogen, C₁-C₅ alkyl, halogen, CHF₂, CH₂F or CF₃;

X is chosen from the group consisting of; -NH-, -O-, -S-, -SO-, -SO₂-, -Se-, -Te- or -S-S-

Y is chosen from the group consisting of; hydrogen, hydroxy, -CH₂OH, methoxy, NH₂, unbranched, branched or cyclic C₁-C₅ alkyl, unbranched, branched or cyclic -NH(C₁-C₅); unbranched, branched or cyclic N(C₁-C₅)₂, -NH(C₆aryl), -N(C₆aryl)₂, -NH(C₁-C₁₀ heteroaryl), and -N(C₅-C₁₀ heteroaryl)₂, C₅-C₁₀ heteroaryl wherein any of said aryl or heteroaryl groups are optionally substituted with up to 3 groups of R^a which groups may be the same or different;

Z is chosen from the group consisting of; C, N, or O;

R^a represents a member selected from: hydrogen, halogen, $-\text{CN}$, OH , CO_2H , CHO , NO_2 , $-\text{NH}_2$, $-\text{NH}(\text{C}_1\text{-C}_4)$, $\text{N}(\text{C}_1\text{-C}_4)_2$, $-\text{NH}(\text{C}_6\text{ aryl})$, $-\text{N}(\text{C}_6\text{ aryl})_2$, $-\text{NH}(\text{C}_5\text{-C}_{10}\text{ heteroaryl})$, and $-\text{N}(\text{C}_5\text{-C}_{10}\text{ heteroaryl})_2$; or a pharmaceutically acceptable salt thereof.

A preferred compound is according to formula I, wherein R_1 or/and R_2 are H, (S)-methyl, methyl, (R)-ethyl, (S)-ethyl, ethyl, (R)-propyl, (S)-propyl, propyl, (S)-butyl, (S)-1-methyl-propyl, (S)-2-methyl-propyl, (R)-isopropyl, (S)-isopropyl, isopropyl, cyclopentyl, $-(\text{CH}_2)_2\text{SMe}$, (R)- $\text{CH}_2\text{SCH}_2\text{Ph}$, (S)-benzyl, 4-chloro-benzyl, (S)-3-methyl-1-H-indole or (S)-phenyl;

Further preferred is a compound according to formula I, wherein R_3 is chosen from the group consisting of; hydrogen, methyl, ethyl, phenyl, 3-hydroxy phenyl, 4-hydroxy phenyl, or forms a keto group together with R_4 .

Further preferred is a compound according to formula I, wherein R_4 is H, methyl, or forms a keto group together with R_3 .

Further preferred is a compound according to formula I, wherein R_5 is NO_2 , CN , CH_2CN or CO_2H ;

Further preferred is a compound according to formula I, wherein R_6 is Me, or CF_3 ;

Further preferred is a compound according to formula I, wherein R_7 is H or Me;

Further preferred is a compound according to formula I, wherein R_8 is H or methyl;

Further preferred is a compound according to formula I, wherein X is NH;


Further preferred is a compound according to formula I, wherein Y is H, -OH, -OMe, -N(CH₂CH₃)₂, piperidine, or 4-nitro-2-ylamino;

Further preferred is a compound according to formula I, wherein Z is CR₇ or N;

Even more preferred is a compound according to formula I, chosen from the group consisting of;










2-Methyl-2-(4-nitro-3-trifluoromethyl-phenylamino)-propan-1-ol;
[1-(4-Nitro-3-trifluoromethyl-phenylamino)-cyclopentyl]-methanol;
(S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-3-phenyl-propan-1-ol;
(S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-butan-1-ol;
2-Methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol;
[1-(3-Methyl-4-nitro-phenylamino)-cyclopentyl]-methanol;
(S)-2-(3-Methyl-4-nitro-phenylamino)-butan-1-ol;
[1-(6-Methyl-5-nitro-pyridin-2-ylamino)-cyclopentyl]-methanol;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-2-phenyl-ethanol;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-3-phenyl-propan-1-ol;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-butan-1-ol;
(DL)-3-(4-Chloro-phenyl)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
(S)-2-(6-Methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
2-(2,3-Dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol;
(S)-2-(3,5-Dimethyl-4-nitro-phenylamino)-butan-1-ol;
4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-benzonitrile;
4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
(S)-4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
(R)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;










(S)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;
[4-((S)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
[4-((R)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
[4-((S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile;
6-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-nicotinonitrile;
4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile;
and the compounds showed in the following table (The substituents, R₉, R₆, and Z, are shown in the table, and are all substituents in the following formula II. In formula II, the NO₂ group corresponds to the substituent R₅ in formula I, and R₉ is composed of the moieties XR₁R₂YR₃R₄ of Formula I as defined above, where X is -NH-






















Formula II

R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH

R9	R6	Z		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		

R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	CH

[illegible]

4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzoic acid;
(6-Methyl-5-nitro-2-pyridin-2-ylamino)-butionic methyl ester;
2-Methyl-N-(6-methyl-5-nitro-pyridin-2-yl amino)-propan-2-ol;
4-((R)-2-Hydroxy-1-methyl-ethylamino)-2-trifluoromethyl-benzonitrile
4-((R)-1-Furan-2-ylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile
(R)-3-Furan-2-yl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol
2-(6-Methyl-5-nitro-pyridin-2-ylamino)-heptan-1-ol
3-Cyclopentyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol
2-(6-Methyl-5-nitro-pyridin-2-ylsulfanyl)-ethanol
[1-(4-Fluoro-3-methyl-phenylamino)-cyclopentyl]-methanol
1-[4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-phenyl]-ethanone
1-[4-((S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-ethanone
1-[4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-phenyl]-ethanone
[1-(4-Methanesulfonyl-3-methyl-phenylamino)-cyclopentyl]-methanol
2,2-Dimethyl-3-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

2, 2-Dimethyl-3-(3-methyl-4-nitro-phenylamino)-propan-1-ol

4-((R)-1-Benzylsulfanylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile

(R)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-3-phenylmethanesulfinyl-propan-1-ol

4-((R)-2-Hydroxy-1-phenylmethanesulfinylmethyl-ethylamino)-2-trifluoromethyl-benzonitrile

[1-(4-Nitro-phenylamino)-cyclopentyl]-methanol

(S)-2-(4-Nitro-phenylamino)-pentan-1-ol

(S)-4-Methyl-2-(4-nitro-phenylamino)-pentan-1-ol

[1-(2-Bromo-4-nitro-phenylamino)-cyclopentyl]-methanol

(S)-2-(2-Bromo-4-nitro-phenylamino)-pentan-1-ol

(S)-2-(2-Bromo-4-nitro-phenylamino)-4-methyl-pentan-1-ol

or a pharmaceutically acceptable salt thereof.

Also preferred is a compound according to Formula I, wherein R_1 or R_2 is a C_6 - C_{10} arylthio moiety comprising an aryl-substituted sulfur-containing C_1 - C_{10} alkyl group.

Further preferred is a compound according to Formula I, wherein in R_1 or R_2 the alkylsulfur is substituted with a C_6 aryl group.

The present invention further provides a pharmaceutical composition which contains one or more of the compounds according to the above.

More preferred is a pharmaceutical composition according to the above, for use as a medicament.

Furthermore, the invention provides the use of a pharmaceutical composition according to the above for manufacturing a medicament to be used in the treatment of a disease caused by a disturbance in the activity of the androgen receptor.

Since the compounds are shown to be mainly antagonists for the androgen receptor, a preferred use is the use of the composition above for treating a disease which is caused by an increase in androgen receptor activity.

Even more preferred is the use of the composition above for treating a disease which is chosen from the group consisting of; prostate cancer, lipid abnormalities, cardiovascular disease and psychological abnormalities, male pattern baldness (alopecia), benign prostatic hyperplasia (BPH) and acne, hirsutism, amenorrhea, hypogonadism, anemia, diabetes, defects in spermatogenesis, cachexia, osteoporosis, osteopenia, and muscle wasting.


The present invention also provides the use of a compound according to the above for manufacturing a medicament to be used in the treatment of a disease caused by a disturbance in the activity of the androgen receptor.

A specific disease that would be amenable for treatment by the present invention is a disease chosen from the group consisting of; prostate cancer, lipid abnormalities, cardiovascular disease and psychological abnormalities, male pattern baldness (alopecia), benign prostatic hyperplasia (BPH) and acne, hirsutism, amenorrhea, hypogonadism, anemia, diabetes, defects in spermatogenesis, cachexia, osteoporosis, osteopenia, and muscle wasting.

Methods of treating such diseases by administering a therapeutically effective amount of such compounds to a patient are also provided by the invention.

The compounds of the present invention can be used alone, in combination with other compounds of the present invention, or in combination with one or more other agent(s) active in the therapeutic areas described herein.

According to another aspect of the invention there is provided a compound as defined in Formula I, provided that the compound is not the compound according to the formula;



The specific compound above is known in the prior art as an intermediate compound in the manufacture of compounds used in different technical fields, namely the dye industry (Compound Reference: Specs and Bio Specs B.V.; Catalog No. AK-079/11126007).

DETAILED DESCRIPTION OF THE INVENTION

The following definitions apply to the terms as used throughout this specification, unless otherwise limited in specific instances.

The term “androgen receptor ligand” as used herein is intended to cover any moiety, which binds to an androgen receptor. The ligand may act as an antagonist, or as a partial antagonist.

A compound being a “partial antagonist” is a compound with both agonistic and antagonistic activity.

The term “alkyl” as employed herein alone or as part of another group refers to an acyclic straight or branched chain radical, containing 1 to about 10 carbons, preferably 1 to 6 carbons in the normal chain, i.e. methyl, ethyl, propyl, isopropyl, sec-butyl, tert-butyl, pentyl, hexyl, octyl. When substituted alkyl is present, this refers to an unbranched or branched alkyl group, which groups may be the same or different at any available point, as defined with respect to each variable.

The term "substituted alkyl" includes an alkyl group optionally substituted with one or more functional groups which are commonly attached to such chains, such as, alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, hydroxy, cyano, nitro, amino, halo, carboxyl or alkyl ester thereof and/or carboxamide.

The term "alkenyl" as employed herein alone or as part of another group refers to a straight or branched chain radical, containing 2 to about 10 carbons, preferably 2 to 6 carbons i.e. ethenyl, propenyl, butenyl, allyl.

The term "allyl" refers to $\text{H}_2\text{C}=\text{CH}-\text{CH}_2$.

The term "alkynyl" as employed herein alone or as part of another group refers to a straight or branched chain radical, containing 2 to about 10 carbons, preferably 2 to 6 carbons i.e. ethynyl, propynyl, butynyl, allyl.

The term "aryl" as employed herein alone or as part of another group refers to substituted and unsubstituted aromatic ring system. The term aryl includes monocyclic aromatic rings, polycyclic aromatic ring system and polyaromatic ring systems. The polycyclic aromatic and polyaromatic ring systems may contain from two to four, more preferably two to three rings. Preferred aryl groups include 5- or 6- membered ring systems.

The term "heteroaryl" refers to optionally substituted aromatic ring system having one or more heteroatoms such as, for example, oxygen, nitrogen and sulfur. The term heteroaryl includes five- or six-membered heterocyclic rings, polycyclic heteroaromatic ring system and polyheteroaromatic ring systems. The poly heterocyclic aromatic and poly heteroaromatic ring systems may contain from two to four, more preferably two to three rings. The term hetero aryl includes ring system such as pyridine, quinoline, furan, thiophene, pyrrole, imidazole and pyrazole.

The term "alkoxy" as employed herein alone or as part of another group refers to an alkyl ether wherein the term alkyl is as defined above. Examples of alkoxy radicals include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy and the like.

The term "aryloxy" as employed herein alone or as part of another group refers to an aryl alkyl ether, wherein the term aryl is as defined above. Examples of aryloxy radicals include phenoxy, benzyloxy and the like.

The term "alkylthio" as employed herein alone or as part of another group refers to an alkyl thio wherein the term alkyl is as defined above and one of the methylene carbons has been replaced with sulfur. Examples of alkylthio radicals include methane thiol, ethane thiol, propane thiol, $-(CH_2)_m S(CH_2)_n$, where $m + n = 9$ and the like.

The term "alkylsulphone" as employed herein alone or as part of another group refers to an alkylsulphone wherein the term alkyl is as defined above and one of the methylene carbons has been replaced with sulfur. Examples of alkylsulphone radicals include methanesulphone, ethanesulphone, propanesulphone, $-(CH_2)_m SO_2(CH_2)_n$, where $m + n = 9$ and the like.

The term "alkylsulphoxide" as employed herein alone or as part of another group refers to an alkylsulphoxide wherein the term alkyl is as defined above and one of the methylene carbons has been replaced with sulfur. Examples of alkylsulphoxide radicals include methanesulphoxide, ethanesulphoxide, propanesulphoxide $-(CH_2)_m SO(CH_2)_n$, where $m + n = 9$ and the like.

The term "alkylarylthio" as employed herein alone or as part of another group refers to an arylalkylthio wherein the term alkylthio and aryl are as defined above and one of the terminal methyl groups is substituted with aryl. Examples of $-(CH_2)_m S(CH_2)_n CH_2-Ar$ where $m + n = 8$ and the like.

The term "alkylarylsulphone" as employed herein alone or as part of another group refers to an arylalkylsulphone wherein the term alkylsulphone and aryl are as defined above and one of the terminal methyl groups is substituted with aryl. Examples of $-(CH_2)_m SO_2(CH_2)_n CH_2-Ar$ where $m + n = 8$ and the like.

The term "alkylarylsulphoxide" as employed herein alone or as part of another group refers to an arylalkylsulphoxide wherein the term alkylsulphoxide and aryl are as defined above and one of the terminal methyl groups is substituted with aryl. Examples of $-(CH_2)_m SO(CH_2)_n CH_2-Ar$ where $m + n = 8$ and the like.

The term "cycloalkyl" as employed herein alone or as part of another group refers to saturated cyclic hydrocarbon groups or partially unsaturated cyclic hydrocarbon groups, independently containing one carbon-to-carbon double bond. The cyclic hydrocarbon contains 3 to 4 carbons. It should also be understood that the present invention also involve cycloalkyl rings where 1 to 2 carbons in the ring are replaced by either -O-, -S- or -N-, thus forming a saturated or partially saturated heterocycle. Examples of such rings

are aziridine, thiiranes and the like. Preferred heterocyclic rings are 3-membered, which may be optionally substituted by 1, 2 or 3 groups of R^a which groups may be the same or different through available carbons as in the case of "alkyl". Preferred cycloalkyl groups include 3 carbons, such as cyclopropyl, which may be optionally substituted by 1, 2 or 3 groups of R^a which groups may be the same or different through available carbons as in the case of "alkyl".

The term "halogen" refers to fluorine, chlorine, bromine and iodine. Also included are carbon substituted halogens such as $-CF_3$, $-CHF_2$, and $-CH_2F$

The compounds of the present invention can be present as salts, which are also within the scope of this invention. Pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts are preferred. If the compounds of the invention have, for example, at least one basic center, they can form acid addition salts. These are formed, for example, with strong inorganic acids, such as mineral acids, for example sulfuric acid, phosphoric acid or a hydrohalic acid, with strong organic carboxylic acids, such as alkanecarboxylic acids of 1 to 4 carbon atoms which are unsubstituted or substituted, for example, by halogen, for example acetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, (for example aspartic or glutamic acid or lysine or arginine), or benzoic acid, or with organic sulfonic acids, such as (C_1 - C_4) alkyl or arylsulfonic acids which are unsubstituted or substituted, for example by halogen, for example methyl- or p-toluene- sulfonic acid. Corresponding acid addition salts can also be formed having, if desired, an additionally present basic center. The compounds of the invention having at least one acid group (e.g. $COOH$) can also form salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine,

piperidine, pyrrolidine, a mono, di or tri-lower alkylamine, for example ethyl, tertbutyl, diethyl, diisopropyl, triethyl, tributyl or dimethyl-propylamine, or a mono, di or trihydroxy lower alkylamine, for example mono, di or triethanolamine. Corresponding internal salts may furthermore be formed. Salts that are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free compounds of the invention or their pharmaceutically acceptable salts, are also included. Preferred salts of the compounds of the present invention which contain a basic group include monohydrochloride, hydrogensulfate, methanesulfonate, phosphate or nitrate. Preferred salts of the compounds of formula I which contain an acid group include sodium, potassium and magnesium salts and pharmaceutically acceptable organic amines.

The compounds according to the invention may also have prodrug forms. Any compound that will be converted in vivo to provide the bioactive agent (i.e., the compound of formula I) is a prodrug within the scope and spirit of the invention. Such prodrugs are well known in the art and a comprehensive description of these may be found in: (i) *The Practice of Medicinal Chemistry*, Camille G. Wermuth et al., Ch 31, (Academic Press, 1996); (ii) *Design of Prodrugs*, edited by H. Bundgaard, (Elsevier, 1985); and (iii) *A Textbook of Drug Design and Development*, P. Krogsgaard-Larson and H. Bundgaard, eds. Ch 5, pgs 113 – 191 (Harwood Academic Publishers, 1991).

Embodiments of prodrugs suitable for use in the present invention include lower alkyl esters, such as ethyl ester, or acyloxyalkyl esters such as pivaloyloxymethyl (POM).

The compounds according to the present invention are preferably administered in a therapeutically effective amount. The term “therapeutically effective amount” as used herein refers to an amount of a therapeutic agent to treat or prevent a condition treatable by administration of a composition of the invention. That amount is the amount sufficient to exhibit a detectable therapeutic or preventative or ameliorative effect. The effect may include, for example, treatment or prevention of the conditions listed herein.


The precise effective amount for a subject will depend upon the subject's size and health, the nature and extent of the condition being treated, recommendations of the treating physician, and the therapeutics or combination of therapeutics selected for administration.

Scheme 1-6 outlines the synthetic routes used for preparing the compound according to Formula I

Scheme 1

Synthetic routes to these compounds can be visualized by the skilled person and the present synthetic route is not limiting for the invention. 4-Fluoro-1-nitro-2-trifluoromethyl-benzene (1a) and 4-fluoro-2-methyl-1-nitro-benzene (1b) were employed as starting material in scheme-1 and is commercially obtainable.

Scheme 1 depicts a synthesis of compounds of formula I in which R_6 is CF_3 and Me and is connected to phenyl ring. Condensation of compound (1a) with different β -amino alcohols and di-isopropyl ethylamine in DMSO gave compound 3 (examples 1-4) in quantitative yield. The reactions were performed in a microwave oven at elevated temperature for a short time. Compound (1b) was used for producing the compound 3 (examples 5-7) and similar conditions were adopted as in examples 1-4. An alternative method was used for the preparation of example-5. The reaction according to the alternative method was performed by heating the compound (1b) and β -amino alcohol in pentanol in a sealed tube.




Scheme 1

Scheme 2

Compounds 9 (examples 8-15) were prepared from starting material 6-chloro-3-nitro-2-picoline (compound 4). Starting material was synthesized in three steps starting with compound 6-amino-2-picoline using the literature procedure. Nitration of 6-amino-2-picoline was accomplished by concentrated sulphuric acid (H_2SO_4) and concentrated nitric acid (HNO_3) and provided 6-amino-3-nitro-2-picoline (Baumgarten, H. E. and Chien Fan Su, H. *JACS* 74 (1952) 3828; Parker, E. D. and Shive, W. *JACS* 69 (1947) 63). Treatment of 6-amino-3-nitro-2-picoline with sodium nitrite provided 6-hydroxy-3-nitro-2-picoline, which, when reacted with PCl_5 and POCl_3 , provided 6-chloro-3-nitro-2-picoline (Baumgarten, H. E. and Chien Fan Su, H. *JACS* 74 (1952) 3828).


Scheme 2 shows the synthesis of compounds of formula I in which Z is N and R_7 is H. Condensation of 6-Chloro-3-nitro-2-picoline and 2-amino-2-methyl-propan-1-ol in 1-pentanol and the mixture refluxed under inert atmosphere gave compound 9 (example-8) as yellow crystals. 6-Chloro-3-nitro-2-picoline can also be purchased as commercial starting material. The reaction time was reduced by using a microwave oven. Condensation of compound 7 with different β -amino alcohols (8) in the microwave provided compound 9 (examples 9-13) in quantitative yield. Synthetic routes to these compounds can be visualized by the skilled person. Reaction of compound (10) with L-alanine provided compound 11 (example-14). Reduction of the acid compound (11) by a reducing agent such as lithium aluminum hydride (LAH) produced compound 9 (example 15).




Scheme 2

Scheme 3


Synthesis of compounds according to formula I, in which R_6 and R_7 are Me and connected to the phenyl ring is shown in Scheme-3. 4-Fluoro-2, 3-di-methyl-1-nitro-benzene (13) was employed as starting material in scheme-3, which was produced by the nitration of compound (12) with fuming nitric acid in acetic anhydride in one step. Condensation of 2, 3-dimethyl-fluoro-benzene with β -amino alcohols at higher temperature gave compound 14 (example 16).

**Scheme 3****Scheme 4**

Scheme 4 depicts a synthesis of compounds of formula I in which R₆ and R₈ are Me and connected to the phenyl ring. Condensation of compound (15) with (S)-2-amino-butan-1-ol and di-isopropyl ethylamine in DMSO gave compound 16 (examples 17). The reaction was performed in a microwave oven.

**Scheme 4****Scheme 5**


Reduction of nitro compound to amine was accomplished by the treatment of sodium thiosulphate with ethanol. After work-up the amines were used for the next step without any further purification. Reaction of amine with potassium cyanide and copper cyanide in water gave compound 19 (examples 26-28). (Clive, D. L. *et al.* *JOC* 52 (1987) 1339-42 and Vogel expt. 6.76). Some other examples of compound 19 were made by an alternative method utilizing a microwave oven. Similar reaction conditions as those used in scheme-1 and scheme-2 provided compound 19 (examples 18-22). Conversion of the nitrile form of compound 19 to benzoic acid compound 20 (example 87) was performed in a refluxed aqueous sodium hydroxide solution in methanol.



Scheme 5

Scheme 6


Scheme 6 depicts a synthesis of compounds of formula I in which R_3 and R_4 are Me and is connected to the alkyl chain. Condensation of 6-chloro-3-nitro-2-picoline with glycine methyl ester hydrochloride and triethyl amine in DMSO gave compound 22 (example 88). Compound 22 was treated with methyl magnesium bromide and after HPLC purification gave compound 23 (example 89).



Scheme 6

EXAMPLES


The following Examples represent preferred embodiments of the present invention. However, they should not be construed as limiting the invention in any way. The ^1H NMR spectra were consistent with the assigned structures. Mass spectra were recorded on a Perkin-Elmer, API 150Ex spectrometer, with turbo "ion spray" on negative ion mode (ES-1) or positive (ES+1), using a Zorbax SB-C8 column (LC-MS). The microwave reactions were performed in a Personal Chemistry Emrys Optimizer.

Example 12-Methyl-2-(4-nitro-3-trifluoromethyl-phenylamino)-propan-1-ol

4-Fluoro-1-nitro-2-trifluoromethyl-benzene (1.226 g, 5.86 mmol) was dissolved in 7 mL DMSO and 2-amino-2-methyl-propan-1-ol (784 mg, 8.795 mmol) was added, followed by diisopropyl ethylamine (DIPEA) (985 mg, 7.622 mmol). The reaction was heated to 180 °C for 900 seconds in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The mixture was diluted with 20 mL of EtOAc and then washed three times with an aqueous solution of ammonium chloride (NH_4Cl). The organic phase was collected, dried with MgSO_4 (anhydrous) and filtered. The dry organic phase was evaporated *in vacuo*. The crude product was a bright yellow powder. The crude product was purified on a silica column with 5:1 n-heptane: EtOAc as mobile

phase. This gave 1.1 g (68 %) of 2-methyl-2-(4-nitro-3-trifluoromethyl-phenylamino)-propan-1-ol as a yellow solid. $M/Z = 278$


Example 2



[1-(4-Nitro-3-trifluoromethyl-phenylamino)-cyclopentyl]-methanol

4-Fluoro-1-nitro-2-trifluoromethyl-benzene (122 mg, 0.583 mmol) was coupled with (1-amino-cyclopentyl)-methanol (101 mg, 0.875 mmol), DIPEA (90.5 mg, 0.700 mmol) in DMSO 0.8 mL, using the same procedure as described in Example-1. This gave 120.5 mg (68%) of [1-(4-nitro-3-trifluoromethyl-phenylamino)-cyclopentyl]-methanol as a yellow powder. $M/Z = 304$.

Example 3




(S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-3-phenyl-propan-1-ol

4-Fluoro-1-nitro-2-trifluoromethyl-benzene (119 mg, 0.569 mmol) was coupled with (S)-2-amino-3-phenyl-propan-1-ol (129 mg, 0.854 mmol), DIPEA (88 mg, 0.683 mmol) in DMSO 0.8 mL using the same procedure as described in Example-1. This gave 112 mg

(58%) of (S)-2-(4-nitro-3-trifluoromethyl-phenylamino)-3-phenyl-propan-1-ol as yellow crystals. $M/Z = 340$.


Example 4



(S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-butan-1-ol

4-Fluoro-1-nitro-2-trifluoromethyl-benzene (122 mg, 0.583 mmol) was coupled with (S)-2-amino-butan-1-ol (78 mg, 0.875 mmol), DIPEA (91 mg, 0.700 mmol) in DMSO 0.8 mL using the same procedure as described in Example-1. This gave 107 mg (67%) of (S)-2-(4-nitro-3-trifluoromethyl-phenylamino)-butan-1-ol as yellow oily crystals. $M/Z = 278$.

Example 5




2-Methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol

Method-A: 4-Fluoro-2-methyl-1-nitro-benzene (113 mg, 0.728 mmol) was coupled with 2-amino-2-methyl-propan-1-ol (84 mg, 0.947 mmol), DIPEA (122 mg, 0.947 mmol) in DMSO 1.2 mL using the same procedure as described in Example-1. The crude product was purified on a silica column with 1:1 n-heptane: EtOAc as mobile phase. This gave 72 mg (44 %) of 2-methyl-2-(3-methyl-4-nitro-phenylamino)-propan-1-ol as yellow powder. $M/Z = 224$.

Method-B: 4-Fluoro-2-methyl-1-nitro-benzene (2.33 g, 15 mmol) and 2-amino-2-methylpropanol (2.67 g, 30 mmol) were heated with stirring at 160°C in a sealed tube overnight. The reaction mixture was diluted with EtOAc and purified by flash chromatography (dry application; 14% EtOAc in hexane → EtOAc) to afford 2.85 g (85%) of the 2-methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol.


Example 6



[1-(3-Methyl-4-nitro-phenylamino)-cyclopentyl]-methanol

4-Fluoro-2-methyl-1-nitro-benzene (107 mg, 0.689 mmol) was coupled with (1-amino-cyclopentyl)-methanol (103 mg, 0.897 mmol), DIPEA (116 mg, 0.897 mmol) in DMSO 1.2 mL using the same procedure as described in Example-1. The crude product was purified on a silica column with 1:1 n-heptane: EtOAc as mobile phase. This gave 76 mg (44 %) of [1-(3-methyl-4-nitro-phenylamino)-cyclopentyl]-methanol as a yellow powder. M/Z = 250.


Example 7



(S)-2-(3-Methyl-4-nitro-phenylamino)-butan-1-ol

4-Fluoro-2-methyl-1-nitro-benzene (102 mg, 0.658 mmol) was coupled with (S)-2-amino-butan-1-ol (76 mg, 0.855 mmol), DIPEA (111 mg, 0.855 mmol) in DMSO 1.2 mL using the same procedure as described in Example-1. The crude product was purified on a silica column with 1:1 n-heptane: EtOAc as mobile phase. This gave 85 mg (58 %) of (S)-2-(3-methyl-4-nitro-phenylamino)-butan-1-ol as yellow oil. M/Z = 224.

Example 8



2-Methyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

(a) Conc. H_2SO_4 (140 ml) was cooled in an ice-salt bath and molten 6-amino-2-picoline (30 g, 0.277 mol) was added in portions with good stirring. To this brown, viscous solution which was maintained at 0°C was added a cooled (0°C) mixture of conc. H_2SO_4 (21 ml) and conc. HNO_3 (21 ml) drop wise over a period of approx. 1.5 hrs. The red-orange reaction mixture was stirred for an additional hour at 0°C and then allowed to warm slowly to room temperature over night. The brown solution was heated at 60°C (oil bath) for 1 hr followed by 1hr at 100°C (carefully controlled temperature). The reaction mixture was cooled to 0°C (ice bath), poured over cracked ice and neutralised by addition of a concentrated aqueous NaOH solution. The yellow precipitate was filtered and washed well with ice-water. (The filtrate was put in the refrigerator; additional product was precipitated together with the salts.) The yellow product was suspended in water and divided into two portions, each of them subjected to steam distillation in turn. The yellow reaction mixture became more "transparent" after some hrs, but the collected steam, containing 4-amino-3-nitro-2-picoline, was still yellow after 6 hrs. The steam distillation was stopped after 8 hrs, the residual part of the reaction mixture was filtered and

evaporated to dryness. ^1H NMR (D_2O) showed a mixture of 2-3 compounds. The mixture was washed with; CHCl_3 , EtOH (x 2) and CHCl_3 leaving 20.4 g (48%) of pure 6-amino-3-nitro-2-picoline.


(b) 6-Amino-3-nitro-2-picoline (20 g, 0.131 mol) was suspended in a mixture of conc. H_2SO_4 (23.7 ml) and water (335 ml). More conc. H_2SO_4 (20 ml) was added under ice-cooling, but the amine did not dissolve completely. The suspension was added in ice (100 g) before a solution of NaNO_2 (13.53 g, 0.196 mol) in water (40 ml) was added drop wise. Gas evolution was observed. The brown suspension was stirred at 10°C for 1 hr, filtered and washed with water. The brown product was dried (freeze dryer) to achieve 15.78 g (78 %) of 6-hydroxy-3-nitro-2-picoline.

(c) To 6-Hydroxy-3-nitro-2-picoline (15.73 g, 0.102 mol) was added PCl_5 (5.73 g, 0.027 mol) and POCl_3 (2.9 ml, 0.032 mol). This mixture was heated at $110\text{--}115^\circ\text{C}$ for 3 hrs. However, the amount of POCl_3 added was only enough to moisten the starting material. More POCl_3 (3 ml) was added, the reaction mixture heated at $110\text{--}115^\circ\text{C}$ but only sublimation of PCl_5 (100°C) was observed. DMF (5 ml) was added and the solution was heated at 115°C for 5 hrs, cooled and poured into a slush of ice and water. A beige product precipitated and the water suspension was stirred for 48 hrs. The brown precipitate was filtered off and washed with water. Purification by dry-flash dichloromethane yielded 10.93 g (62 %) of 6-chloro-3-nitro-2-picoline.

(d) 6-Chloro-3-nitro-2-picoline (6.055 g, 35.1 mmol) and 2-amino-2-methyl-propan-1-ol (6.2 g, 73.7 mmol) were suspended in 1-pentanol (30 ml) and the mixture refluxed under inert atmosphere overnight. The thin layer chromatography (dichloromethane 4/EtOAc 1) revealed some remaining starting material, so the reaction was refluxed for another 3.5 hrs. The reaction mixture was cooled and water was added under stirring. A sticky, yellow precipitate was filtered off, washed well with water and dried. The crude product (6.04 g) was re-crystallised from either pentane-acetone or dichloromethane. Collecting

the crops furnished 5.71 (72 %) of 2-methyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol as yellow crystals. $M/Z = 225$.


Example 9



[1-(6-Methyl-5-nitro-pyridin-2-ylamino)-cyclopentyl]-methanol

6-Chloro-3-nitro-2-picoline (22 mg, 0.13 mmol) was coupled with (1-amino-cyclopentyl)-methanol (31 mg, 0.27 mmol), triethylamine (0.025 mL, 0.18 mmol) in 2-pentanol (1 mL). The reaction was heated to 180 °C for 2 h in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The mixture was diluted with 20 mL of EtOAc and then washed with NaHCO₃. The organic phase was collected, dried with anhydrous MgSO₄ and filtered. The dry organic phase was evaporated and purified on a silica column with 5:1 n-Heptane: EtOAc as mobile phase. This gave 9 mg (28%) of [1-(6-methyl-5-nitro-pyridin-2-ylamino)-cyclopentyl]-methanol as a yellow solid. $M/Z = 251$

Example 10




(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-2-phenyl-ethanol

6-Chloro-3-nitro-2-picoline (22 mg, 0.13 mmol) was coupled with (2-amino-2-phenyl)-


propanol (34 mg, 0.25 mmol) in triethylamine (0.030 mL, 0.25 mmol) in DMSO (1 mL). The reaction was heated to 140 °C for 1200 seconds in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The mixture was diluted with 20 mL of EtOAc and then washed with NH₄Cl (aq) three times. The organic phase was collected, dried with anhydrous MgSO₄ and filtered. The dry organic phase was evaporated and purification on silica column with 5:1 n-Heptane: EtOAc gave 22 mg (63%) of (R)-2-(6-methyl-5-nitro-pyridin-2-ylamino) 2-phenyl-ethanol as a yellow solid. M/Z = 273.

Example 11




(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-3-phenyl-propan-1-ol.

6-Chloro-3-nitro-2-picoline (30 mg, 0.17 mmol) was coupled with (S)-2-amino-3-phenyl-propan-1-ol (32 mg, 0.21 mmol), sodium acetate (28 mg, 0.34 mmol) in EtOH (2 mL). The reaction was heated in a microwave oven for 20 min at 130 °C and then additionally 20 minutes at 150 °C. The reaction was quenched with a saturated aqueous solution of NaHCO₃ and extracted with EtOAc and evaporated. Purification on a silica column with a gradient solution of heptane: EtOAc gave 24 mg (48%) of (S)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-3-phenyl-propan-1-ol as a yellow solid. M/Z = 287.

Example 12


(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-butan-1-ol

6-Chloro-3-nitro-2-picoline (30 mg, 0.17 mmol) was coupled with (S)-2-amino-butan-1-ol (32 mg, 0.21 mmol), and sodium acetate (28 mg, 0.34 mmol) in EtOH (2 mL) using the same procedure as described in Example-13. This gave 21 mg (53%) of (S)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-butan-1-ol as a yellow solid. $M/Z = 225$.


Example 13

(DL)-3-(4-Chloro-phenyl)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol.

6-Chloro-3-nitro-2-picoline (50 mg, 0.29 mmol) was coupled with (DL)-2-amino-3-(4-chloro-phenyl)-propan-1-ol (103 mg, 0.55 mmol), in triethylamine (0.077 mL, 0.55 mmol) in DMSO (1 mL) using the same procedure as described in Example-1 but at 140 °C. This gave 23 mg (45%) of (DL)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-3-(4-chloro-phenyl)-propan-1-ol as a yellow solid. $M/Z = 321$.

Example 14**(S)-2-(6-Methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid**


6-Chloro-3-nitro-2-picoline (62 mg, 0.36 mmol) was coupled with L-alanine (80 mg, 0.90 mmol) and sodium acetate (78 mg, 0.95 mmol) in DMSO 1 mL. The reaction was heated to 140 °C for 600 seconds in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The crude mixture was treated with a saturated aqueous solution of NH_4Cl . The reaction mixture was acidified to pH 4 (HCl , 1M). The crude reaction mixture was extracted with EtOAc, and the combined organic layers were washed with water and brine. Purification on silica using a mobile phase CH_2Cl_2 -MeOH-HOAc gave 60 mg (74%) of (S)-2-(6-methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid as a yellow solid. $M/Z = 225$.

Example 15**(S)-2-(6-Methyl-5-nitro-2-pyridin-2-ylamino)-propan-1-ol**

(S)-2-(6-Methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid (60 mg, 0.27 mmol) was added to a nitrogen-purged flask with LiAlH_4 (27 mg, 0.71 mmol). The reaction mixture was refluxed for 2 h and then allowed to reach room temperature and then quenched by

sequentially adding H₂O (1 mL), NaOH (1M, 1 mL) and H₂O (1 mL). The slurry was centrifuged and the precipitated aluminum salts were washed with dichloromethane. The combined filtrates were evaporated and purification of the residue on a silica column with heptane- EtOAc (3:2) gave 13 mg (22%) of (S)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol as a yellow solid. M/Z = 211.

Example 16




2-(2,3-Dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol

Fuming nitric acid (1.4 g, 20.3 mmol) was cooled to 0°C and acetic anhydride (2.89 g, 28.4 mmol) was added. This solution was added to a cold (0°C) solution of 3-fluoro-1,2-dimethylbenzene (1.0 g, 8.1 mmol) in acetic anhydride (4 ml) over 10 min. The reaction mixture was stirred for 25 min, poured slowly over ice and the water solution extracted with EtOAc (x 3). The collected organic phase was washed with diluted saturated aqueous solution of NaHCO₃ followed by brine before evaporation to dryness. The residue was flash purified on a silica gel column using hexane as a mobile phase to give 2,3-dimethyl-4-fluoro-1-nitro-benzene 0.74 g (54%) as a yellow oil which crystallised upon standing.

The fluoride (0.576 g, 3.4 mmol) was mixed with 2-amino-2-methylpropanol (0.61 g, 6.8 mmol) in a tube, and the tube was sealed before immersing it into an oil bath and heating at 160°C for 5 days. TLC (Hexane) showed remaining starting material. The reaction mixture was cooled and diluted with EtOAc before purification by flash silica gel chromatography (dry application; 6:4 hexane and EtOAc) to give 0.34 g (59% recovery) of the starting material 2,3-dimethyl-4-fluoro-1-nitro-benzene and 0.20 g (61% based on

recovered starting material) of the 2-(2,3-dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol. $M/Z = 238$.


Example 17



(S)-2-(3,5-Dimethyl-4-nitro-phenylamino)-butan-1-ol

(S)-2-Amino-butan-1-ol (41 mg, 0.461 mmol) was dissolved in DMSO (800 μ L) and DIPEA (80 μ L, 0.461 mmol) added. 4-Fluoro-2-trifluoromethyl-benzonitrile (60mg, 0.354 mmol) was added and the reaction mixture was heated to 160 $^{\circ}$ C for 900 seconds in a microwave oven (Parameters: High absorbance, Fixed Holding time, pre-stirring 25 sec). The reaction mixture was then diluted with EtOAc and washed with an aqueous solution of NH_4Cl . The organic phase was then dried and evaporated *in vacuo*. The crude product was purified on silica column with 3:1 n-heptane:EtOAc as the mobile phase. This provided 22 mg (26 %) of (S)-2-(3,5-dimethyl-4-nitro-phenylamino)-butan-1-ol. $M/Z = 238$


Example 18



4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-benzonitrile


2-Amino-2-methyl-propan-1-ol (25 mg, 0.275 mmol) was dissolved in 0.7 mL DMSO and DIPEA (36 mg, 0.275 mmol) was added. 4-fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.212 mmol) was then added and the reaction was heated to 140 °C for 1100 seconds in a microwave oven (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The reaction was then diluted with 10 mL EtOAc, washed with an aqueous solution of NH₄Cl, dried with anhydrous MgSO₄, filtered and then the organic phase was evaporated *in vacuo*. The crude product was purified on silica column with 3:1 n-heptane:EtOAc as the mobile phase. Upon dissolving the crude product in the mobile phase, an insoluble precipitate was collected. On analysis this showed to be mainly pure product. All insoluble precipitate was dissolved in acetone, celite™ was added, whereafter the acetone was evaporated. The celite was then applied to a silica column with 2:1 n-heptane:EtOAc as the mobile phase to give 34 mg (62%) of 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-benzonitrile as beige crystals. M/Z = 258.

Example 19




4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile

4-Fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.212 mmol) was coupled with (1-amino-cyclopentyl)-methanol (32 mg, 0.275 mmol), and DIPEA (36 mg, 0.275 mmol) in DMSO 0.7 mL using the same procedure as described in Example-8. This gave 23 mg (38%) of 4-(1-hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile as white powder. M/Z = 284.

Example 20


(S)-4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile

4-Fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.212 mmol) was coupled with (S)-2-amino-butan-1-ol (25 mg, 0.275 mmol), DIPEA (36 mg, 0.275 mmol), in 0.7 mL DMSO using the same procedure as described in Example-8. This gave 17 mg (31%) of (S)-4-(1-hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile as white crystals. $M/Z = 258$.

Example 21


(R)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile

4-Fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.21 mmol), (R)-2-Amino-pentan-1-ol (32 mg, 0.27 mmol) and DIPEA (47 μ L, 0.27 mmol) was dissolved in DMSO (1 mL) and heated to 180 °C for 900 seconds in a microwave oven (Parameters: Fixed Holding time, High absorbance, pre-stirring 25 sec.). The crude product was diluted with CH_2Cl_2 and washed with an aqueous solution of NH_4Cl . The organic phase was separated, dried and evaporated in vacuo. The crude product was purified on a silica column with 3:1 n-heptane: EtOAc as the mobile phase. This gave 39 mg (68%) of (R)-4-(1-hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile. $M/Z = 272$.

Example 22

(S)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile

4-Fluoro-2-trifluoromethyl-benzonitrile (40 mg, 0.21 mmol) was coupled with (S)-2-Amino-pentan-1-ol (32 mg, 0.27 mmol), DIPEA (47 μ L, 0.27 mmol) in DMSO 1.0 mL, using the same procedure as described in Example-21. This gave 24 mg (42%) of (S)-4-(1-hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile. M/Z = 272


Example 23

[4-(R)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile

(4-Fluoro-2-trifluoromethyl-phenyl)-acetonitrile (100 mg, 0.492 mmol) was dissolved in DMSO (3.5 mL) and (R)-(-)-2-Amino-1-pentanol (66 mg, 0.634 mmol) and pyridine (52 μ L, 0.634 mmol) was added. The reaction was heated in microwave to 170 $^{\circ}$ C for 900 sec (Parameters: 30 seconds pre-stirring, holding time on, normal absorption). The mixture was diluted with EtOAc and washed with aqueous solution of NH_4Ac . The water phase was washed with EtOAc and the organic phases were pooled, dried with MgSO_4 , filtered

and evaporated *in vacuo*. The crude product was purified on a silica column with 5:1 n-heptane: EtOAc as the mobile phase. This gave 2.1 mg (1.5 %) of [4-(R)-1-hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile. $M/Z = 286$


Example 24



[4-(S)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile

(4-Fluoro-2-trifluoromethyl-phenyl)-acetonitrile (100 mg, 0.492 mmol) was coupled with (S)-(+)-2-Amino-1-pentanol (66 mg, 0.634 mmol), Pyridine (52 μ L, 0.634 mmol), in DMSO (3.5 mL) using the same procedure as described in Example-23. This gave 2.2 mg (1.6 %) of [4-(S)-1-hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile. $M/Z = 286$

Example 25




[4-(S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile

(4-Fluoro-2-trifluoromethyl-phenyl)-acetonitrile (119 mg, 0.584 mmol) was coupled with L-Leucinol (89 mg, 0.759 mmol), Pyridine (62 μ L, 0.759 mmol), DMSO (3.2 mL) using

the same procedure as described in Example-23. This gave 2.6 mg (1.5 %) of [4-((S)-1-hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile. M/Z = 300

Example 26




4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile

The 2-methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol (360 mg, 1.6 mmol) was dissolved in ethanol (26 ml) and $\text{Na}_2\text{S}_2\text{O}_4$ (2.23 g, 12.8 mmol) was added and the solution heated at 80°C overnight. The solvent was evaporated and the remaining solid was partitioned between 10% aqueous solution NaHCO_3 and EtOAc. The water phase (pH = neutral) was extracted with EtOAc (x 3), the collected organic phase washed with brine and dried (MgSO_4). The 2-(4-amino-3-methyl-phenylamino)-2-methyl-propan-1-ol was used in the next step without further purification. (The amine oxidises on the TLC plate; brown spots upon standing.)

Sodium nitrite (NaNO_2) (190 mg, 2.75 mmol) in water (2.5 ml) was added to a solution of amine (500 mg, 2.5 mmol conc. HCl /ice (2.5 ml/2.5 g) during 5 min. followed by neutralisation by addition of solid CaCO_3 . KCN (391 mg, 6 mmol) and CuCN (269 mg, 3.0 mmol) in water (1 ml) was heated at 60°C (oil bath) and the cold, neutral diazonium salt solution was added drop wise over 15 min. Gas evolution was observed and the resulting suspension turned bright and strong orange. The reaction mixture was heated at 110°C for 30 min, cooled, diluted with water and EtOAc and filtered through celite. The water phase was extracted with EtOAc and the collected organic phase washed with brine and dried (MgSO_4). The crude product (491 mg) was purified by flash chromatography

(Hexane; Hex/EtOAc; 7:3 → 1:1) giving the reduced compound 2-methyl-2-(3-hydroxy-phenylamino)-propan-1-ol (93 mg) and 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile (108 mg, 21%) as a pale yellow solid. $M/Z = 204$.

Example 27




6-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-nicotinonitrile

2-Methyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol (1.08 g, 4.8 mmol) was dissolved in 75% aqueous ethanol and $\text{Na}_2\text{S}_2\text{O}_4$ (3.9 g, 24 mmol) was added in portions. The reaction mixture was heated at 60°C for 30 min when TLC (10% MeOH in DCM) showed full conversion. The heat was turned off, the reaction mixture stirred overnight at ambient temperature and evaporated to dryness. To this residue was added NaHCO_3 (5% aq.) and EtOAc, the phases separated and the water phase (pH 7-8) extracted extensively with EtOAc. (The product is very water-soluble and it is probably better to do a continuous extraction with EtOAc to get a higher yield). The collected organic phase was washed with brine before drying (MgSO_4). Upon standing, the colour of the organic solution turned from yellow to orange. Filtration and evaporation yielded 0.648 g (69%) of amine as a red oil.

NaNO_2 (0.25 g, 3.65 mmol) in water (3 ml) was added to a solution of amine 6 (0.648 g, 3.3 mmol) in ice/conc. HCl (3.5 g/3.5 ml) during 5 min. followed by neutralisation by addition of solid CaCO_3 . KCN (0.52 g, 7.96 mmol) and CuCN (0.36 g, 3.98 mmol) in water (3 ml) was heated at 60°C (oil bath) and the cold, neutral diazoniumsalt solution was added drop wise over 15 min. Gas evolution was observed and the resulting

suspension turned bright and strong orange. The reaction mixture was heated at 110°C for 30 min, cooled, diluted with water and EtOAc and filtered through celite. The water phase was extracted with EtOAc and the collected organic phase was washed with brine and dried (MgSO₄). The crude product (0.248 g) was purified by flash chromatography (Hexane → Hex :EtOAc 3:7) yielding 34 mg of 2-methyl-2-(6-methyl-pyridin-2-ylamino)-propan-1-ol and 11 mg of 6-(2-hydroxy-1,1-dimethyl-ethylamino)-2-methyl-nicotinonitrile. $M/Z = 205$.

Example 28



4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile

The nitro compound 18 (0.20 g, 0.84 mmol) was dissolved in EtOH (20 ml), Na₂S₂O₄ (1.1 g, 6.71 mmol) was added and the reaction mixture heated at 80°C overnight. The cold reaction mixture was filtered through celite, washed well with EtOAc and the filtrate evaporated to dryness. The crude 2-(4-amino-2,3-dimethyl-phenylamino)-2-methyl-propan-1-ol (0.292 g), pure by ¹H-NMR, was used as such in the next reactions.

The reaction was performed using the same procedure as described in Example-21 using 2-(4-amino-2,3-dimethyl-phenylamino)-2-methyl-propan-1-ol (0.175 g, 0.84 mmol) in conc. HCl/ice water (1 ml/5 ml), NaNO₂ (64 mg, 0.92 mmol) in water (1 ml), KCN (130 mg, 2 mmol) and CuCN (90 mg, 1 mmol) in water (1 ml). The crude product (341 mg) was purified by flash chromatography (Hexane; Hex 7/EtOAc 3) giving reduced compound 2-(2,3-dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol and 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile. All the fractions containing impure nitrile were collected and crystallised from hexane/EtOAc to give 25

mg (13%) of pure 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile.
M/Z = 218.


Procedure for Library synthesis (Examples 29-86).

The following is the general procedure for library synthesis for the examples of 29-88.
The compounds are shown in table 2.









Smith-vials for the microwave oven were charged with 0.1 mmol either of the starting materials; 5-fluoro-2-nitro toluene, 5-fluoro-2-nitrobenzotrifluoride, 6-fluoro-2-methyl-3-nitro-pyridine.










To each vial was added 0.5 ml DMSO, 20 μ L triethylamine (1.4 equivalents), and 1.4 equivalents of the diverse amino alcohols. The vials were run 1100s in 140°C in a microwave oven. After synthesis the products were analysed by LC-MS. The DMSO solutions were transferred to test tubes, and evaporated onto silica gel under reduced pressure. The silica gel from the tubes was placed on SPE SI columns, and a frit was placed on top. The products were purified with a gradient solution of heptane/EtOAc. The fractions were pooled and solvent was evaporated. Compounds which were more than 90% pure were tested in an *in vitro* assay which is described below. Purity was determined by analytic HPLC.










The scaffold used for the construction of the library is according to Formula II. The



Formula II












Example	R9	R6	Z	Yield (%)	MS (-Q1)
29		CF ₃	CH	46	262.9
30		CF ₃	CH	55	290.8
31		CF ₃	CH	24	249.1
32		CF ₃	CH	62	276.7
33		CF ₃	CH	65	290.8
34		CF ₃	CH	23	290.8
35		CF ₃	CH	93	325.3
36		CF ₃	CH	78	341.2

Example	R9	R6	Z	Yield (%)	MS (-Q1)
37		CF ₃	CH	82	262.9
38		CF ₃	CH	95	305.2
39		CF ₃	CH	98	323.2
40		CF ₃	CH	98	290.8
41		CF ₃	CH	89	385
42		CF ₃	CH	92	290.8
43		CF ₃	CH	95	290.8
44		CF ₃	CH	100	378.1
45		CF ₃	CH	84	316

Example	R9	R6	Z	Yield (%)	MS (-Q1)
46		CF ₃	CH	90	262.9
47		CF ₃	CH	106	275.2
48		CF ₃	CH	75	304.3
49		CF ₃	CH	69	275.2
50		CF ₃	CH	76	370
51		CF ₃	CH	89	325.3
52		CH ₃	N	53	238.0
53		CH ₃	N	53	238.0
54		CH ₃	N	30	195.7

Example	R9	R6	Z	Yield (%)	MS (-Q1)
55		CH ₃	N	60	223.9
56		CH ₃	N	63	238.0
57		CH ₃	N	22	238.0
58		CH ₃	N	88	272.2
59		CH ₃	N	65	209.8
60		CH ₃	N	60	252.1
61		CH ₃	N	79	252.1
62		CH ₃	N	89	252.1
63		CH ₃	N	74	270.4

Example	R9	R6	Z	Yield (%)	MS (-Q1)
64		CH ₃	N	84	238.0
65		CH ₃	N	78	332.2
66		CH ₃	N	88	238.0
67		CH ₃	N	80	224.2
68		CH ₃	N	75	238.0
69		CH ₃	N	72	209.8
70		CH ₃	N	58	223.1
71		CH ₃	N	52	222.1
72		CH ₃	N	90	272.2
73		CH ₃	C	44.0	208.9

Example	R9	R6	Z	Yield (%)	MS (-Q1)
74		CH ₃	CH	55.0	237.1
75		CH ₃	CH	66.0	195.1
76		CH ₃	CH	31.0	237.1
77		CH ₃	CH	30.0	223
78		CH ₃	CH	32.0	237.1
79		CH ₃	CH	27	271.3
80		CH ₃	CH	25	250.9
81		CH ₃	CH	27	269.2
82		CH ₃	CH	24	237.1
83		CH ₃	CH	24	237.1
84		CH ₃	CH	24	237.1





Example	R9	R6	Z	Yield (%)	MS (-Q1)
85		CH ₃	CH	25	250
86		CH ₃	CH	33	316

Table 2


Example 87**4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzoic acid**

A suspension of 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile (70 mg, 0.34 mmol) and NaOH (0.14 g, 3.4 mmol) in water/MeOH (5 ml/8 ml) was refluxed for 4 days. The reaction mixture was diluted with water, pH adjusted to approx. 3 with 50% aq. HCl. The precipitated solid was filtered off and collected, the water phase was extracted with EtOAc (x 3), washed with brine and dried (MgSO₄). The crude product was purified on a silica column with 1:1 n-heptane: EtOAc as mobile phase. This gave 39 mg (51%) of the 4-(2-hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzoic acid as a brownish foam. M/Z 223.

Example-88


(6-Methyl-5-nitro-2-pyridin-2-ylamino)-butionic methyl ester

6-chloro-3-nitro-2-picoline (600 mg, 3.5 mmol) was coupled with glycine methyl ester hydrochloride (880 mg, 7 mmol), triethylamine (1.5 ml, 10.5 mmol) in DMSO 3 mL at 140 °C for 30 min in microwave (Parameters: high absorbance, fixed holding time, pre-stirring 25 seconds). The crude mixture was treated with a saturated aqueous solution of NH_4Cl . The aqueous solution of was extracted with EtOAc, washed with water and brine. The crude product was purified on a silica column with CH_2Cl_2 -MeOH as mobile phase. This gave 39 mg (51%) of 580 mg (74%) of (6-methyl-5-nitro-2-pyridin-2-ylamino)-butionic methyl ester as a yellow solid. $M/Z = 225$.

Example-89


2-Methyl-N-(6-methyl-5-nitro-pyridin-2-yl amino)-propan-2-ol

2-(6-Methyl-5-nitro-pyridin-2-ylamino)-butionic methyl ester (30 mg, 0.13 mmol) was dissolved in THF (3 mL) and added to a nitrogen-purged flask containing methyl magnesium chloride (MeMgCl) (0.08 ml, 0.027 mmol) at 0 °C. The reaction mixture was allowed to reach room temperature and then refluxed for 5 h. The reaction was quenched by adding saturated NH_4Cl . The reaction mixture was extracted with EtOAc and washed with H_2O and brine. The crude product was purified by HPLC. This gave 1.5 mg (5%) of 2-methyl-N-(6-methyl-5-nitro-pyridin-2-yl amino)-propan-2-ol as yellow oil. $M/Z = 225$.

Example-90

4-((R)-2-Hydroxy-1-methyl-ethylamino)-2-trifluoromethyl-benzonitrile


D-Alanine (36 mg, 0.40 mmol) was dissolved in THF (dry, 1 ml) and the vials were purged with N₂ for 5 min. BF₃-Et₂O (0.050 ml 0.40 mmol) was added with syringe and the mixture was heated at 70°C for 1.5 h. BH₃-SMe₂ (0.22 ml, 0.44 mmol, 2M solution) was added carefully during vigorous stirring (an exotherm was formed approx half way) (a evolution of gas was noticed). The reactions was purged with N₂ and then heated at 70°C over night (17h). The reaction was allowed to cool to room temp. The excess borane was quenched by addition of 1 ml of a 1:1 mixture of THF: H₂O, followed by 1 ml of NaOH (5M). The two phase system was heated at 70°C in 4h. The flask was purged with N₂ to blow off the THF. CH₂Cl₂ (2 ml) was added and the two phase system was transformed to a Phase separator. Additional CH₂Cl₂ (2 ml) was added and the combined organic phases were evaporated. The crude (21 mg) was then dissolved in DMSO and the reaction was continued as in example 1. 4-Fluoro-2-trifluoromethyl-benzonitrile (19 mg, 0.1 mmol) was coupled with the formed (R)-2-amino-propan-1-ol. DIPEA (0.021 ml, 0.12 mmol), in 1 mL DMSO using the same procedure as described in Example-1. Purification on preperative HPLC gave 4 mg (16 %) of 4-((R)-2-Hydroxy-1-methyl-ethylamino)-2-trifluoromethyl-benzonitrile as a white solid. M/Z= 244.

Example-91

4-((R)-1-Furan-2-ylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile

(R)-2-Amino-3-furan-2-yl-propionic acid (40 mg, 0.25 mmol) was reduced using the same procedure as described in Example-90. The crude was coupled with 4-Fluoro-2-trifluoromethyl-benzonitrile (19 mg, 0.1 mmol) and DIPEA (0.05 ml, 0.2 mmol) as in example 1 and gave 4-((R)-1-furan-2-ylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile 11 mg (29%), after purification on HPLC, as a white solid. $M/Z = 310$.


Example-92



(R)-3-Furan-2-yl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

(R)-2-Amino-3-furan-2-yl-propionic acid (40 mg, 0.25 mmol) was reduced using the same procedure as described in Example-90. The crude was coupled with 6-chloro-3-nitro-2-picoline (17 mg, 0.1 mmol) and DIPEA (0.05 ml, 0.2 mmol) as in example 1 and gave, after purification on HPLC, 9 mg (33%) of (R)-3-Furan-2-yl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol, as a white solid. $M/Z = 277$.

Example-93




2-(6-Methyl-5-nitro-pyridin-2-ylamino)-heptan-1-ol

2-Amino-heptanoic acid (33 mg, 0.25 mmol) was reduced using the same procedure as described in Example-90. The crude was coupled with 6-chloro-3-nitro-2-picoline (17 mg, 0.1 mmol) and DIPEA (0.05 ml, 0.2 mmol) as in Example 1 and gave after

purification on HPLC, 3 mg (11 %) 2-(6-methyl-5-nitro-pyridin-2-ylamino)-heptan-1-ol, as an oil. M/Z= 267


Example-94



3-Cyclopentyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

2-Amino-3-cyclopentyl-propionic acid (36 mg, 0.25 mmol) was reduced using the same procedure as described in Example-90. The crude was coupled with 6-chloro-3-nitro-2-picoline (17 mg, 0.1 mmol) and DIPEA (0.05 ml, 0.2 mmol) as in Example 1 and gave, after purification on HPLC, 4 mg (14 %) 3-Cyclopentyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol, as a yellow solid. M/Z= 279.


Example-95



2-(6-Methyl-5-nitro-pyridin-2-ylsulfanyl)-ethanol

6-Chloro-3-nitro-2-picoline (17 mg, 0.1 mmol) was coupled with 2-Mercapto-ethanol (0.014 ml, 0.2 mmol), DIPEA (25 mg, 0.2 mmol) in DMSO 0.8 mL, using the same procedure as described in Example-1. This gave 5 mg (23%) of 2-(6-Methyl-5-nitro-pyridin-2-ylsulfanyl)-ethanol as a yellow oil. M/Z = 214.


Example 96



[1-(4-Fluoro-3-methyl-phenylamino)-cyclopentyl]-methanol

4-Fluoro-2-methyl phenol (0.24 mmol) was solved in 800 μL DMSO. (1-Amino-cyclopentyl)-methanol (0.29 mmol) was added and then Diisopropyl-ethyl amine (DIPEA) (0.29 mmol). Reaction was heated to 180 $^{\circ}\text{C}$ in microwave for 15 min (Parameters: Normal absorption, hold time on, pre-stirring 20 sec). Starting material was remaining so reaction was heated to 220 $^{\circ}\text{C}$ for additional 15 min. Several products obtained. Crude mixture was diluted in CH_2Cl_2 and washed several times with NH_4Cl (aq) and phases were separated on SPE Phase Separator. Organic phase was evaporated *in vacuo* and crude product mixture was then purified on silica column with 5:1 n-heptane:EtOAc as mobile phase. This gave 2.3 mg (4 %) of [1-(4-fluoro-3-methyl-phenylamino)-cyclopentyl]-methanol. $M/Z = 221$


Example 97



1-[4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-phenyl]-ethanone


1-(4-Fluoro-2-trifluoromethyl-phenyl)-ethanone (40mg, 0.194 mmol) was solved in 800 μL DMSO. 2-Amino-2-methyl-propan-1-ol (23mg, 0.252 mmol) was added and then DIPEA (44 μL , 0.252 mmol). Reaction mixture was heated to 180 $^{\circ}\text{C}$ in microwave for 15 min (Parameters: Normal absorption, hold time on, pre-stirring 25 sec). Majority of starting material still left so reheated to 210 $^{\circ}\text{C}$ for 15 min. Several products obtained. Crude mixture was diluted in CH_2Cl_2 and washed several times with NH_4Cl (aq) and phases were separated on SPE Phase Separator. Organic phase was evaporated *in vacuo* and crude product mixture was then purified on silica column with 10:1 n-heptane:EtOAc as mobile phase. This gave 3 mg (6%) of 1-[4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-phenyl]-ethanone as minor product. $M/Z = 275$.

Example 98




1-[4-((S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-ethanone

1-(4-Fluoro-2-trifluoromethyl-phenyl)-ethanone (40 mg, 0.194 mmol) was coupled with (S)-2-Amino-4-methyl-pentan-1-ol (30mg, 0.252 mmol), DIPEA (44 μL , 0.252 mmol) in DMSO 800 μL using the same procedure as described in Example-97. This gave 15 mg (25 %) of 1-[4-((S)-1-hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-ethanone. $M/Z = 303$

Example 99


1-[4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-phenyl]-ethanone

1-(4-Fluoro-2-trifluoromethyl-phenyl)-ethanone (40 mg, 0.194 mmol) was coupled with (1-Amino-cyclopentyl)-methanol (29 mg, 0.252 mmol), DIPEA (44 μ L, 0.252 mmol), in DMSO 800 μ L using the same procedure as described in Example-97. This gave 5 mg (9 %) of 1-[4-(1-hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-phenyl]-ethanone. M/Z = 301.

Example 100

[1-(4-Methanesulfonyl-3-methyl-phenylamino)-cyclopentyl]-methanol

4-Fluoro-1-methanesulfonyl-2-methyl-benzene (40 mg, 0.213 mmol) was solved in 800 μ L DMSO. (1-Amino-cyclopentyl)-methanol (32 mg, 0.276 mmol) was added and DIPEA (48 μ L, 0.276 mmol). Reaction mixture was heated to 180 $^{\circ}$ C in microwave for 15 min (Parameters: Normal absorption, hold time on, pre-stirring 30 sec). Crude mixture was diluted in CH_2Cl_2 and washed several times with NH_4Cl (aq) and phases were separated on SPE Phase Separator. Organic phase was evaporated *in vacuo* and crude product mixture was then purified on silica column with 7:1 n-heptane:EtOAc as mobile phase. This gave 1.4 mg (2 %) of [1-(4-methanesulfonyl-3-methyl-phenylamino)-cyclopentyl]-methanol. M/Z = 283


Example 101

2,2-Dimethyl-3-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

6-Chloro-2-methyl-3-nitro-pyridine (40 mg, 0.232 mmol) was solved in 900 μ L DMSO. 3-Amino-2,2-dimethyl-propan-1-ol (31 mg, 0.301 mmol) and DIPEA (52 μ L, 0.301 mmol) was added and heated to 180 $^{\circ}$ C in microwave for 15 min parameters: Normal

absorption, hold time on, pre-stirring 30 sec). Crude mixture was diluted in CH_2Cl_2 and washed several times with NH_4Cl (aq) and phases were separated on SPE Phase Separator. Organic phase was evaporated *in vacuo* and crude product mixture was then purified on silica column with 7:1 n-heptane:EtOAc as mobile phase. This gave 15 mg (27 %) of 2,2-dimethyl-3-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol. $M/Z = 239$


Example 102



2, 2-Dimethyl-3-(3-methyl-4-nitro-phenylamino)-propan-1-ol

4-Fluoro-2-methyl-1-nitro-benzene (40 mg, 0.258 mmol) was coupled with 3-Amino-2,2-dimethyl-propan-1-ol (35 mg, 0.335 mmol), DIPEA (58 μL , 0.335 mmol) in DMSO 900 μL using the same procedure as described in Example-101. This gave 4 mg (7 %) of 2, 2-dimethyl-3-(3-methyl-4-nitro-phenylamino)-propan-1-ol. $M/Z = 238$.


Example 103



4-((R)-1-Benzylsulfanylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile

4-Fluoro-2-trifluoromethyl-benzonitrile (60 mg, 0.32 mmol) was solved in 1000 μL DMSO. (R)-2-amino-3-benzylsulfanyl-propan-1-ol (81 mg, 0.41 mmol) was added and then diisopropyl-ethyl amine (DIPEA) (53 mg, 0.41 mmol). Reaction was heated to 180 $^{\circ}\text{C}$ in microwave for 15 min (Parameters: Normal absorption, hold time on, pre-stirring 20 sec). Crude mixture was diluted in CH_2Cl_2 and washed several times with NH_4Cl (aq) and phases were separated on SPE Phase Separator. Organic phase was evaporated *in vacuo* and crude product mixture was then purified on silica column with 3:1 n-heptane:EtOAc as mobile phase. This gave pure product 82 mg (71 %) of 4-((R)-1-benzylsulfanylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile as transparent oil. $M/Z = 366$


Example 104



(R)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-3-phenylmethanesulfinyl-propan-1-ol

CH₂Cl₂ (0.125 mL) was cooled to 0 °C and mCPBA (13 mg, 0.07 mmol) was solved in it. Stirred at 0 °C for 10 min then (R)-3-benzylsulfonyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol (20 mg, 0.06 mmol) was added. Stirred at 0 °C for 20 min. Cooling bath was removed and reaction was allowed to warm to room temperature and was then stirred overnight. The organic phase was washed with brine, phases were separated on SPE Phase Separator and organic phase was dried and evaporated *in vacuo*. Crude product gives precipitation on salivation in 3:1 n-Heptane:EtOAc. Precipitate was consisting of mainly product and was solved in acetonitrile and purified on silica column with EtOAc as mobilephase. This gave 8.2 mg (39 %) of (R)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-3-phenylmethanesulfinyl-propan-1-ol. M/Z = 349.


Example 105



4-((R)-2-Hydroxy-1-phenylmethanesulfinylmethyl-ethylamino)-2-trifluoromethyl-benzonitrile

4-((R)-1-Benzylsulfonylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile (20 mg, 0.06 mmol) was reacted with mCPBA (11 mg, 0.07 mmol) in CH₂Cl₂ (0.125 mL) using the same procedure as described in Example-104. This gave 14.1 mg (67 %) of 4-((R)-2-Hydroxy-1-phenylmethanesulfinylmethyl-ethylamino)-2-trifluoromethyl-nicotinonitrile. M/Z = 382


Example 106



[1-(4-Nitro-phenylamino)-cyclopentyl]-methanol

1-Fluoro-4-nitro-benzene (41 mg, 0.29 mmol) was solved in 1000 μ L DMSO. (1-amino-cyclopentyl)-methanol (44 mg, 0.38 mmol) was added and then diisopropyl-ethyl amine (DIPEA) (49 mg, 0.38 mmol). Reaction was heated to 170 $^{\circ}$ C in microwave for 15 min (Parameters: Normal absorption, hold time on, pre-stirring 30 sec). Crude mixture was diluted in EtOAc and washed several times with NH_4Cl (aq) and phases were separated. Organic phase was dried and then evaporated *in vacuo*. Crude product mixture was purified on silica column with 3:1 n-heptane:EtOAc as mobile phase. This gave 48 mg (70 %) of [1-(4-nitro-phenylamino)-cyclopentyl]-methanol. $M/Z = 236$.


Example 107



(S)-2-(4-Nitro-phenylamino)-pentan-1-ol

1-Fluoro-4-nitro-benzene (41 mg, 0.29 mmol) was coupled with (S)-2-amino-pentan-1-ol (39 mg, 0.38 mmol), diisopropyl-ethyl amine (DIPEA) (49 mg, 0.38 mmol) in DMSO 1000 μ L using the same procedure as described in Example-106. This gave 53 mg (81 %) of (S)-2-(4-nitro-phenylamino)-pentan-1-ol. $M/Z = 224$


Example-108



(S)-4-Methyl-2-(4-nitro-phenylamino)-pentan-1-ol

1-Fluoro-4-nitro-benzene (42 mg, 0.29 mmol) was coupled with (S)-2-amino-4-methyl-pentan-1-ol (50 mg, 0.38 mmol), diisopropyl-ethyl amine (DIPEA) (50 mg, 0.38 mmol) in DMSO 1000 μ L using the same procedure as described in Example-106. This gave 40 mg (57 %) of (S)-4-Methyl-2-(4-nitro-phenylamino)-pentan-1-ol. $M/Z = 238$.


Example 109



[1-(2-Bromo-4-nitro-phenylamino)-cyclopentyl]-methanol

[1-(4-Nitro-phenylamino)-cyclopentyl]-methanol (10 mg, 0.042 mmol) was solved in a 1:1 mixture of CH_2Cl_2 : MeOH (2 mL). CaCO_3 (8.5 mg, 0.085 mmol) was added and the solution was stirred at roomtemp for 10 min. Benzyltrimethylammonium tribromide (36 mg, 0.093 mmol) was added and the reaction was stirred at roomtemp for 48 h. Crude reaction was diluted with CH_2Cl_2 and washed with $\text{NH}_4\text{Cl}_{(\text{aq})}$. Organic phase was collected, dried and evaporated *in vacuo*. Crude product was purified on silica column. This gave 11 mg (83 %) of [1-(2-bromo-4-nitro-phenylamino)-cyclopentyl]-methanol. $M/Z = 315$.


Example 110



(S)-2-(2-Bromo-4-nitro-phenylamino)-pentan-1-ol

(S)-2-(4-Nitro-phenylamino)-pentan-1-ol (29 mg, 0.13 mmol) was treated benzyltrimethylammonium tribromide (111 mg, 0.29 mmol) and CaCO_3 (26 mg, 0.26 mmol) in 1:1 mixture of CH_2Cl_2 : MeOH (2 mL) using the same procedure as described in Example-109. This gave 16 mg (41 %) of (S)-2-(2-Bromo-4-nitro-phenylamino)-pentan-1-ol. $M/Z = 303$.

Example 111



(S)-2-(2-Bromo-4-nitro-phenylamino)-4-methyl-pentan-1-ol

(S)-4-Methyl-2-(4-nitro-phenylamino)-pentan-1-ol (29 mg, 0.13 mmol) was treated benzyltrimethylammonium tribromide (111 mg, 0.30 mmol) and CaCO_3 (26 mg, 0.27 mmol) in 1:1 mixture of CH_2Cl_2 :MeOH (2 mL) using the same procedure as described in Example-109. This gave 20 mg (47 %) of (S)-2-(2-Bromo-4-nitro-phenylamino)-4-methyl-pentan-1-ol. $M/Z = 317$.

All molecules were named by Autonom 2000, part of was IS/Draw 2.5

All naming done by was IS/Draw 2.5 with Autonom 2000

Example- 112

AR Competition Binding Assay

Recombinant human androgen receptor (hAR) was extracted from Sf9 insect cells with buffer containing 1 mM EDTA, 20 mM K_2HPO_4 , 8.7% glycerol, 20 mM Na_2MoO_4 and 12 mM MTG at 5×10^7 cells/ml. The cell debris was removed by centrifugation and the supernatant aliquoted and stored at $-70^\circ C$.

An aliquot of AR extract was thawed on ice prior to use and diluted to approximately 0.2 nM (1 to 30 dilution) in buffer (100 mM $K_nH_mPO_4$ pH 7.0, 1 mM EDTA, 8.7% glycerol, 20 mM Na_2MoO_4 and 1 mM DTT). The test ligands were diluted in DMSO as a dilution series of 10 concentrations in duplicate, with 1:5 dilution between each concentration. Tritiated mibolerone (3H -Mib) was used as tracer compound and diluted to 1.6 nM in 1 mM EDTA, 20 mM Na_2MoO_4 , 8.7% glycerol and 1 mM DTT. To a 96-well polypropylene-plate 110 μ l/well of 1.6 nM 3H -Mib, 10 μ l/well test substance and 110 μ l/well diluted AR was added. The plates were covered and incubated at $+4^\circ C$ over night. The plates were harvested on filters to separate bound ligand from unbound ligand with a Tomtec Harvester. A prewet buffer containing 20 mM $K_n(PO_4)$ pH 7.6, 1 mM EDTA, v/v 0.5% polyethyleneimine was used to equilibrate the filter before filtering the samples and washing the filters with 20 mM $K_n(PO_4)$ pH 7.6, 1 mM EDTA 8 times. The filters were allowed to dry for 1 hour at $+65^\circ C$. A scintillating wax was melted upon the filter and the radioactivity retained on the filter was measured in a Wallac Microbeta scintillation counter.

The affinity to AR was evaluated by a non-linear four-parameter logistic model: $b = (b_{\max} - b_{\min}) / (1 + (IC_{50}/I)^S) + b_{\min}$, where b_{\max} = total concentration of binding sites, b_{\min} = non-specific binding, I = added concentration of binding inhibitor, IC_{50} = concentration of binding inhibitor at half-maximal binding and S = slope factor.

Table: Antagonist and partial antagonist and binding activity of androgen receptor modulator compounds.

AR Transactivation Assays

The agonist and antagonist properties of compounds were determined using a cell-based system expressing stably integrated androgen receptor and an androgen responsive reporter gene. CV-1 cells (kidney fibroblasts) stably expressing CMV-hAR and alkaline phosphatase (ALP) driven by an MMTV promoter containing an androgen response element were cultured in Dulbecco's Modified Eagle Medium (DMEM), low glucose supplemented with 10% fetal bovine serum, 1% L-glutamine, and 0.7% Hygromycin B. The stably integrated cells (ARAF) were trypsinized and resuspended in Opti-MEM 1 supplemented with 2% fetal bovine serum, 1% L-Glutamine, 50 µg/ml Gentamicin and 1% Pen/Strep. The cells were counted in a Birch chamber and diluted to a concentration of 100 000 cells /ml. The cells were then seeded out in 384 plates, 5000cells/well in 50µl seeding media and incubated overnight in 37 C, 5% CO₂.

The next day, the seeding medium was removed from the cells and 20 µl induction media (Opti-MEM 1 supplemented with 1% L- Glutamine, 50 µg/ml Gentamicin and 1% Pen/Strep) +/- 0.1 nM Mibolerone was added to the wells. 10µl of test compound diluted in induction media was then added to the wells. The cells were incubated 48 hr in 37 C, 5% CO₂.

After 48 hr 5µl of cell medium was added to white 384 plates with 100µl of ALP substrate buffer. The plates were incubated in 37 C for 20 minutes followed by incubation at room temperature for 10 minutes before each well was read in a µBETA machine. Agonist activity was calculated from the alkaline phosphatase activity induced


in the absence of Mibolerone and compared to standard activation curve generated by Mibolerone alone. Antagonist activity was calculated from the decrease in ALP activity in the presence of 0.1 nM Mibolerone. EC50 and IC50 values were calculated by using a non-linear four-parameter fit as described above.

Other assays to determine androgen receptor mediated activity of the test compounds include modulation of endogenous AR mediated transcription in cell culture systems; modulation of androgen responsive tissue effects in rodents; identification of receptor surface conformation changes; and binding specificity to AR versus other nuclear receptors.

	AR_LT IC50 (nM)	ARAF EC50 (nM)	ARAF %AGONIST	ARAF IC50 (nM)	ARAF % ANTAGONIST
Example-1	22.77	26.8	51.7	2.1	33.1
Example-5	38.06	81.7	29.3	7.2	61
Example-8	241.44	374.2	10.6	22.3	82.5
Example-19	130.38			22.4	95.6
Example-30	113.45	1069.9	7.3	68.3	88.7
Example-41	65.10			490.3	71.7
Example-42	485.50			493.3	92
Example-60	68.30	336.3	9.3	27.4	79.3
Example-61	89.30			68.3	87.4
Example-65	6.20	1867.3	7	78.0	89.2
Example-78	54.50	279.7	25	25.8	65.4
Example-86	443.40			350.7	100
Example-107	98.70			135.5	92.9
Example-110	170.30			240.7	86.2

CLAIMS

1. Use of a compound according to Formula I in the manufacture of a medicament for the treatment of a disease caused by a disturbance in the activity of the androgen receptor, wherein Formula I is defined as:



Formula I

in which;

R₁ and R₂ are the same or different and independently selected from the group consisting of; hydrogen, halogen, C₁-C₁₀ alkyl, C₁-C₁₀ substituted alkyl, C₂-C₁₀ alkenyl, C₂-C₁₀ alkynyl, C₁-C₁₀ alkoxy, C₁-C₁₀ alkenoxy, C₁-C₁₀ alkynoxy, C₁-C₁₀ alkylthio, C₁-C₁₀ alkenylthio, C₁-C₁₀ alkynylthio, C₆-C₁₀ arylthio, C₁-C₁₀ alkylsulphone, C₁-C₁₀ alkenylsulphone, C₁-C₁₀ alkynylsulphone, C₆-C₁₀ arylsulphone, C₁-C₁₀ alkylsulphoxide, C₁-C₁₀ alkenylsulphoxide, C₁-C₁₀ alkynylsulphoxide, C₆-C₁₀ arylsulphoxide, C₁-C₁₀ alkylarylthio, C₁-C₁₀ alkylarylulphone, C₁-C₁₀ alkylarylulphoxide, C₆-C₁₀ aryl, or C₅-C₂₀ heteroaryl, optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be the same or different; or R₁ and R₂ may together form a C₃-C₁₀ cycloalkyl group;

R₃ and R₄ are the same or different and independently selected from hydrogen, halogen, C₁-C₂₀ alkyl, C₃-C₇ cycloalkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ alkenoxy, C₁-C₄ alkynoxy, C₁-C₄ alkylthio, C₁-C₄ alkenylthio, C₁-C₄ alkynylthio, C₁-C₁₀ alkylsulphone, C₁-C₁₀ alkenylsulphone, C₁-C₁₀ alkynylsulphone, C₆-C₁₀ arylsulphone, C₁-C₁₀ alkylsulphoxide, C₁-C₁₀ alkenylsulphoxide, C₁-C₁₀ alkynylsulphoxide, C₆-C₁₀ arylsulphoxide, C₁-C₁₀ alkylarylthio, C₁-C₁₀ alkylarylulphone, C₁-C₁₀ alkylarylulphoxide, C₆-C₁₅ aryl, C₅-C₂₀ heteroaryl optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be the same or different; or can together form a keto group;

R₅ is chosen from the group consisting of; nitro, cyano, -CH₂CN, -COMe, acetic acid, halogen, sulphonic acid, -SO₂CH₃, aldehyde, carboxylic acid or ester, phosphonic acid or ester;

R₆ is chosen from the group consisting of; hydrogen, C₁-C₅ alkyl, halogen, CN, CO₂H, CHF₂, CH₂F or CF₃;

R₇ is chosen from the group consisting of; H, halogen or C₁-C₅ alkyl;

R₈ is chosen from the group consisting of; hydrogen, C₁-C₅ alkyl, halogen, CHF₂, CH₂F or CF₃;

X is chosen from the group consisting of; -NH-, -O-, -S-, -SO-, -SO₂-, -Se-, -Te- or -S-S-


Y is chosen from the group consisting of; hydrogen, hydroxy, -CH₂OH, methoxy, NH₂, unbranched, branched or cyclic C₁-C₅ alkyl, unbranched, branched or cyclic -NH(C₁-C₈); unbranched, branched or cyclic N(C₁-C₈)₂, -NH(C₆aryl), -N(C₆aryl)₂, -NH(C₁-C₁₀ heteroaryl), and -N(C₅-C₁₀ heteroaryl)₂, C₅-C₁₀ heteroaryl wherein any of said aryl or heteroaryl groups are optionally substituted with up to 3 groups of R^a which groups may be the same or different;

Z is chosen from the group consisting of; C, N, or O;









R^a represents a member selected from: hydrogen, halogen, -CN, OH, CO₂H, CHO, NO₂, -NH₂, -NH(C₁-C₄); N(C₁-C₄)₂, -NH(C₆aryl), -N(C₆aryl)₂, -NH(C₅-C₁₀ heteroaryl), and -N(C₅-C₁₀ heteroaryl)₂; or a pharmaceutically acceptable salt thereof.










2. Use according to claim 1, wherein R_1 or/and R_2 are H, (S)-methyl, methyl, (R)-ethyl, (S)-ethyl, ethyl, (R)-propyl, (S)-propyl, propyl, (S)-butyl, (S)-1-methyl-propyl, (S)-2-methyl-propyl, (R)-isopropyl, (S)-isopropyl, isopropyl, cyclopentyl, $-(CH_2)_2SMe$, (R)- CH_2SCH_2Ph , (S)-benzyl, 4-chloro-benzyl, (S)-3-methyl-1-H-indole or (S)-phenyl;
3. Use according to either of the preceding claims wherein R_3 is chosen from the group consisting of; hydrogen, methyl, ethyl, phenyl, 3-hydroxy phenyl, 4-hydroxy phenyl, or forms a keto group together with R_4 .
4. Use according to any of the preceding claims wherein R_4 is H, methyl, or forms a keto group together with R_3 .
5. Use according to any of the preceding claims wherein R_5 is NO_2 , CN, CH_2CN or CO_2H ;
6. Use according to any of the preceding claims wherein R_6 is Me, or CF_3 ;
7. Use according to any of the preceding claims wherein R_7 is H or Me;
8. Use according to any of the preceding claims wherein R_8 is H or methyl;
9. Use according to any of the preceding claims wherein X is NH;
10. Use according to any of the preceding claims wherein Y is H, -OH, -OMe, -N $(CH_2CH_3)_2$, piperidine, or 4-nitro-2-ylamino;
11. Use according to any of the preceding claims wherein Z is CR_7 or N;
12. Use according to any of the preceding claims wherein the compound is chosen from the group consisting of;
 - 2-Methyl-2-(4-nitro-3-trifluoromethyl-phenylamino)-propan-1-ol;
 - [1-(4-Nitro-3-trifluoromethyl-phenylamino)-cyclopentyl]-methanol;
 - (S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-3-phenyl-propan-1-ol;










(S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-butan-1-ol;
2-Methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol;
[1-(3-Methyl-4-nitro-phenylamino)-cyclopentyl]-methanol;
(S)-2-(3-Methyl-4-nitro-phenylamino)-butan-1-ol;
2-Methyl-2-(6-methyl-5-nitro-pyridine-2-ylamino)-propan-1-ol;
[1-(6-Methyl-5-nitro-pyridine-2-ylamino)-cyclopentyl]-methanol;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino) 2-phenyl-ethanol;
(S)-2-(6-Methyl-5-nitro-pyridine-2-ylamino)-3-phenyl-propan-1-ol;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-butan-1-ol;
(DL)-3-(4-Chloro-phenyl)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
(S)-2-(6-Methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid;
(S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
2-(2,3-Dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol;
(S)-2-(3,5-Dimethyl-4-nitro-phenylamino)-butan-1-ol;
4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-benzonitrile;
4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
(S)-4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
(R)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;
(S)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;
[4-((S)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
[4-((R)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
[4-((S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile;
6-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-nicotinonitrile;
4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile;
and compounds having the formula:













in which R_9 , R_6 and Z are as defined in the following table:












R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH

R9	R6	Z		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		

R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	CH

R9	R6	Z		
	CH ₃	CH		
	CH ₃	CH		
	CH ₃	CH		
	CH ₃	CH		
	CH ₃	CH		
	CH ₃	CH		
	CH ₃	CH		
	CH ₃	CH		
	CH ₃	CH		
	CH ₃	CH		
	CH ₃	CH		

[illegible]

4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzoic acid;

(6-Methyl-5-nitro-2-pyridin-2-ylamino)-butionic methyl ester,

2-Methyl-N-(6-methyl-5-nitro-pyridin-2-yl amino)-propan-2-ol;

4-((R)-2-Hydroxy-1-methyl-ethylamino)-2-trifluoromethyl-benzonitrile

4-((R)-1-Furan-2-ylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile

(R)-3-Furan-2-yl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

2-(6-Methyl-5-nitro-pyridin-2-ylamino)-heptan-1-ol

3-Cyclopentyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

2-(6-Methyl-5-nitro-pyridin-2-ylsulfanyl)-ethanol

[1-(4-Fluoro-3-methyl-phenylamino)-cyclopentyl]-methanol

1- [4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-phenyl]-ethanone

1-[4-((S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-ethanone

1-[4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-phenyl]-ethanone

[1-(4-Methanesulfonyl-3-methyl-phenylamino)-cyclopentyl]-methanol

2,2-Dimethyl-3-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

2,2-Dimethyl-3-(3-methyl-4-nitro-phenylamino)-propan-1-ol

4-((R)-1-Benzylsulfanylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile

(R)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-3-phenylmethanesulfinyl-propan-1-ol

4-((R)-2-Hydroxy-1-phenylmethanesulfinylmethyl-ethylamino)-2-trifluoromethyl-benzonitrile

[1-(4-Nitro-phenylamino)-cyclopentyl]-methanol

(S)-2-(4-Nitro-phenylamino)-pentan-1-ol

(S)-4-Methyl-2-(4-nitro-phenylamino)-pentan-1-ol

[1-(2-Bromo-4-nitro-phenylamino)-cyclopentyl]-methanol

(S)-2-(2-Bromo-4-nitro-phenylamino)-pentan-1-ol

(S)-2-(2-Bromo-4-nitro-phenylamino)-4-methyl-pentan-1-ol


or a pharmaceutically acceptable salt thereof.

13. Use of compound according to claim 1, wherein R_1 or R_2 is a C_6 - C_{10} arylthio comprising an aryl-substituted sulfur-containing C_1 - C_{10} alkyl group.

14. Use of a compound according to claim 1, wherein in R_1 or R_2 the alkylsulfur is substituted with a C_6 aryl group.

15. A pharmaceutical composition containing a compound as defined in Formula I of any preceding claim.

16. Use according to claim 1 wherein the disease is caused by an increase in androgen receptor activity.
17. Use according to any of claims 1-14 or 16 wherein the disease is chosen from the group consisting of, prostate cancer, lipid abnormalities, cardiovascular disease and psychological abnormalities, male pattern baldness (alopecia), benign prostatic hyperplasia (BPH) and acne, hirsutism, amenorrhea, hypogonadism, anemia, diabetes, defects in spermatogenesis, cachexia, osteoporosis, osteopenia, and muscle wasting.
18. A compound as defined by Formula I :



Formula I

in which;

R_1 and R_2 are the same or different and independently selected from the group consisting of; hydrogen, halogen, C_1 - C_{10} alkyl, C_1 - C_{10} substituted alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_1 - C_{10} alkoxy, C_1 - C_{10} alkenoxy, C_1 - C_{10} alkynoxy, C_1 - C_{10} alkylthio, C_1 - C_{10} alkenylthio, C_1 - C_{10} alkynylthio, C_6 - C_{10} arylthio, C_1 - C_{10} alkylsulphone, C_1 - C_{10} alkenylsulphone, C_1 - C_{10} alkynylsulphone, C_6 - C_{10} arylsulphone, C_1 - C_{10} alkylsulphoxide, C_1 - C_{10} alkenylsulphoxide, C_1 - C_{10} alkynylsulphoxide, C_6 - C_{10} arylsulphoxide, C_1 - C_{10} alkylarylthio, C_1 - C_{10} alkylarylsulphone, C_1 - C_{10} alkylarylsulphoxide, C_6 - C_{10} aryl, or C_3 - C_{20} heteroaryl, optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be the same or different; or R_1 and R_2 may together form a C_3 - C_{10} cycloalkyl group;

R_3 and R_4 are the same or different and independently selected from hydrogen, halogen, C_1 - C_{20} alkyl, C_3 - C_7 cycloalkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, C_1 - C_4 alkoxy, C_1 - C_4 alkenoxy, C_1 - C_4 alkynoxy, C_1 - C_4 alkylthio, C_1 - C_4 alkenylthio, C_1 - C_4 alkynylthio, C_1 - C_{10} alkylsulphone, C_1 - C_{10} alkenylsulphone, C_1 - C_{10} alkynylsulphone, C_6 - C_{10} arylsulphone, C_1 - C_{10} alkylsulphoxide, C_1 - C_{10} alkenylsulphoxide, C_1 - C_{10} alkynylsulphoxide, C_6 - C_{10} arylsulphoxide, C_1 - C_{10} alkylarylthio, C_1 - C_{10} alkylarylsulphone, C_1 - C_{10}

alkylarylsulphoxide, C₆-C₁₅ aryl, C₅-C₂₀ heteroaryl optionally substituted with 0, 1, 2 or 3 groups of R^a which groups may be the same or different; or can together form a keto group;

R₅ is chosen from the group consisting of; nitro, cyano, -CH₂CN, -COMe, acetic acid, halogen, sulphonic acid, -SO₂CH₃, aldehyde, carboxylic acid or ester, phosphonic acid or ester;

R₆ is chosen from the group consisting of; hydrogen, C₁-C₅ alkyl, halogen, CN, CO₂H, CHF₂, CH₂F or CF₃;

R₇ is chosen from the group consisting of; H, halogen or C₁-C₅ alkyl;

R₈ is chosen from the group consisting of; hydrogen, C₁-C₅ alkyl, halogen, CHF₂, CH₂F or CF₃;


X is chosen from the group consisting of; -NH-, -O-, -S-, -SO-, -SO₂-, -Se-, -Te- or -S-S-

Y is chosen from the group consisting of; hydrogen, hydroxy, -CH₂OH, methoxy, NH₂, unbranched, branched or cyclic C₁-C₅ alkyl, unbranched, branched or cyclic -NH(C₁-C₅); unbranched, branched or cyclic N(C₁-C₅)₂, -NH(C₆aryl), -N(C₆aryl)₂, -NH(C₁-C₁₀ heteroaryl), and -N(C₅-C₁₀ heteroaryl)₂, C₅-C₁₀ heteroaryl wherein any of said aryl or heteroaryl groups are optionally substituted with up to 3 groups of R^a which groups may be the same or different;

Z is chosen from the group consisting of; C, N, or O;


R^a represents a member selected from: hydrogen, halogen, -CN, OH, CO₂H, CHO, NO₂, -NH₂, -NH(C₁-C₄); N(C₁-C₄)₂, -NH(C₆aryl), -N(C₆aryl)₂, -NH(C₅-C₁₀ heteroaryl), and -N(C₅-C₁₀ heteroaryl)₂; or a pharmaceutically acceptable salt thereof.

with the proviso that the compound is not:


































































































































19. A compound according to claim 18, wherein R_1 or/and R_2 are H, (S)-methyl, methyl, (R)-ethyl, (S)-ethyl, ethyl, (R)-propyl, (S)-propyl, propyl, (S)-butyl, (S)-1-methyl-propyl, (S)-2-methyl-propyl, (R)-isopropyl, (S)-isopropyl, isopropyl, cyclopentyl, $-(CH_2)_2SMe$, (R)- CH_2SCH_2Ph , (S)-benzyl, 4-chloro-benzyl, (S)-3-methyl-1-H-indole or (S)-phenyl;
20. A compound according to either of claims 18 and 19, wherein R_3 is chosen from the group consisting of; hydrogen, methyl, ethyl, phenyl, 3-hydroxy phenyl, 4-hydroxy phenyl, or forms a keto group together with R_4 .
21. A compound according to any of claims 18-20, wherein R_4 is H, methyl, or forms a keto group together with R_3 .
22. A compound according to any of claim 18-21, wherein R_5 is NO_2 , CN, CH_2CN or CO_2H ;
23. A compound according to any of claims 18-22, wherein R_6 is Me, or CF_3 .
24. A compound according to any of claims 18-23, wherein R_7 is H or Me.
25. A compound according to any of claims 18-24, wherein R_8 is H or methyl.
26. A compound according to any of claims 18-25, wherein X is NH.
27. A compound according to any of claims 18-26, wherein Y is H, -OH, -OMe, -N $(CH_2CH_3)_2$, piperidine, or 4-nitro-2-ylamino.
28. A compound according to any of claims 18-27, wherein Z is CR_7 or N.
29. A compound according to any of claims 18-28, wherein the compound is chosen from the group consisting of:










2-Methyl-2-(4-nitro-3-trifluoromethyl-phenylamino)-propan-1-ol;
 [1-(4-Nitro-3-trifluoromethyl-phenylamino)-cyclopentyl]-methanol;
 (S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-3-phenyl-propan-1-ol;
 (S)-2-(4-Nitro-3-trifluoromethyl-phenylamino)-butan-1-ol;
 2-Methyl-2-(3-hydroxy-4-nitro-phenylamino)-propan-1-ol;
 [1-(3-Methyl-4-nitro-phenylamino)-cyclopentyl]-methanol;
 (S)-2-(3-Methyl-4-nitro-phenylamino)-butan-1-ol;
 2-Methyl-2-(6-methyl-5-nitro-pyridine-2-ylamino)-propan-1-ol;
 [1-(6-Methyl-5-nitro-pyridine-2-ylamino)-cyclopentyl]-methanol;
 (S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-2-phenyl-ethanol;
 (S)-2-(6-Methyl-5-nitro-pyridine-2-ylamino)-3-phenyl-propan-1-ol;
 (S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-butan-1-ol;
 (DL)-3-(4-Chloro-phenyl)-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
 (S)-2-(6-Methyl-5-nitro-2-pyridin-2-ylamino)-propionic acid;
 (S)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol;
 2-(2,3-Dimethyl-4-nitro-phenylamino)-2-methyl-propan-1-ol;
 (S)-2-(3,5-Dimethyl-4-nitro-phenylamino)-butan-1-ol;
 4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-benzonitrile;
 4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
 (S)-4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-benzonitrile;
 (R)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;
 (S)-4-(1-Hydroxymethyl-butylamino)-2-trifluoromethyl-benzonitrile;
 [4-((S)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
 [4-((R)-1-Hydroxymethyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
 [4-((S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-acetonitrile;
 4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzonitrile;
 6-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-nicotinonitrile;
 4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2,3-dimethyl-benzonitrile;
 and compounds having the formula:






















in which R₉, R₆ and Z are as defined in the following table:

R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH

R9	R6	Z		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃	CH		
	CF ₃			

R9	R6	Z
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CF ₃	CH
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N

R9	R6	Z
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	N
	CH ₃	CH

[illegible]

[illegible]

4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-methyl-benzoic acid;

(6-Methyl-5-nitro-2-pyridin-2-ylamino)-butionic methyl ester,

2-Methyl-N-(6-methyl-5-nitro-pyridin-2-yl amino)-propan-2-ol;

4-((R)-2-Hydroxy-1-methyl-ethylamino)-2-trifluoromethyl-benzonitrile

4-((R)-1-Furan-2-ylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile

(R)-3-Furan-2-yl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

2-(6-Methyl-5-nitro-pyridin-2-ylamino)-heptan-1-ol

3-Cyclopentyl-2-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

2-(6-Methyl-5-nitro-pyridin-2-ylsulfanyl)-ethanol

[1-(4-Fluoro-3-methyl-phenylamino)-cyclopentyl]-methanol

1-[4-(2-Hydroxy-1,1-dimethyl-ethylamino)-2-trifluoromethyl-phenyl]-ethanone

1-[4-((S)-1-Hydroxymethyl-3-methyl-butylamino)-2-trifluoromethyl-phenyl]-ethanone

1-[4-(1-Hydroxymethyl-cyclopentylamino)-2-trifluoromethyl-phenyl]-ethanone

[1-(4-Methanesulfonyl-3-methyl-phenylamino)-cyclopentyl]-methanol

2,2-Dimethyl-3-(6-methyl-5-nitro-pyridin-2-ylamino)-propan-1-ol

2,2-Dimethyl-3-(3-methyl-4-nitro-phenylamino)-propan-1-ol

4-((R)-1-Benzylsulfanylmethyl-2-hydroxy-ethylamino)-2-trifluoromethyl-benzonitrile

(R)-2-(6-Methyl-5-nitro-pyridin-2-ylamino)-3-phenylmethanesulfinyl-propan-1-ol

4-((R)-2-Hydroxy-1-phenylmethanesulfinylmethyl-ethylamino)-2-trifluoromethyl-benzonitrile

[1-(4-Nitro-phenylamino)-cyclopentyl]-methanol

(S)-2-(4-Nitro-phenylamino)-pentan-1-ol

(S)-4-Methyl-2-(4-nitro-phenylamino)-pentan-1-ol

[1-(2-Bromo-4-nitro-phenylamino)-cyclopentyl]-methanol

(S)-2-(2-Bromo-4-nitro-phenylamino)-pentan-1-ol

(S)-2-(2-Bromo-4-nitro-phenylamino)-4-methyl-pentan-1-ol

30. A compound according to any of claims 18-29, wherein R₁ or R₂ is a C₆-C₁₀ arylthio comprising an aryl-substituted sulfur-containing C₁-C₁₀ alkyl group.

31. A compound according to any of claims 18-30, wherein in R₁ or R₂ the alkylsulfur is substituted with a C₆ aryl group.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB2004/004464

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C205/11 C07C255/50 C07C323/25 C07D213/74 A61K31/04
A61K31/435 A61K31/277 A61K31/10 A61P5/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, BEILSTEIN Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01/58854 A (BIOPHYSICA, INC) 16 August 2001 (2001-08-16) page 4, line 22 - page 5, line 3 claim 1	1-31
A	WO 02/16310 A (GTX, INC; DALTON, JAMES; MILLER, DUANE, D; YIN, DONGHUA; HE, YALI) 28 February 2002 (2002-02-28) claims 12,14,16,18,26 ----- -/--	1-31

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

9 March 2005

Date of mailing of the international search report

18/03/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Goetz, G

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/GB2004/004464

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; KELLER, HELMUT ET AL: "Oxidative hair dye composition comprising a 2,5-diaminobenzonitrile" XP002320584 retrieved from STN Database accession no. 127:311348 see RN 197382-91-5 abstract & EP 0 797 980 A1 (WELLA A.-G., GERMANY) 1 October 1997 (1997-10-01)</p>	18-28
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; HASHIZUME, KAZUNARI ET AL: "Oxidizable color-producing reagents containing p-fluoroaniline derivatives" XP002320585 retrieved from STN Database accession no. 117:127839 see RN 143205-22-5 abstract & EP 0 488 756 A1 (WAKO PURE CHEMICAL INDUSTRIES, LTD., JAPAN) 3 June 1992 (1992-06-03)</p>	18-21
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; TEACH, EUGENE G.: "Herbicidal oxazolidines and methods of use" XP002320586 retrieved from STN Database accession no. 108:217796 see RN 114010-11-6 abstract & US 4 723 986 A (TEACH, EUGENE G.) 9 February 1988 (1988-02-09)</p>	18-21
X	<p>DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SUDA, HIDEAKI ET AL: "p-Nitroaniline derivatives" XP002320587 retrieved from STN Database accession no. 81:25336 see RN 52177-12-5 abstract & JP 49 020129 A2 (SUMITOMO CHEMICAL CO., LTD.) 22 February 1974 (1974-02-22)</p>	18-28
	-/--	

INTERNATIONAL SEARCH REPORT

Inte 1al Application No
PCT/GB2004/004464

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p> DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ZORINA, L. N. ET AL: "New approach to the synthesis of N-aryl-1,3-oxazolidines and N-aryl-1,3-tetrahydrooxazines" XP002320588 retrieved from STN Database accession no. 112:216820 see RN 126993-10-0 abstract & DOKLADY AKADEMII NAUK SSSR , 308(5), 1150-4 'CHEM.! CODEN: DANKAS; ISSN: 0002-3264, 1989, ----- </p>	18-21

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB2004/004464

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0158854	A	16-08-2001	US 6472415 B1	29-10-2002
			AT 259781 T	15-03-2004
			AU 3685601 A	20-08-2001
			AU 5302600 A	20-08-2001
			BR 0008439 A	23-04-2002
			CA 2400185 A1	16-08-2001
			CN 1416416 A	07-05-2003
			CZ 20012398 A3	14-11-2001
			DE 10084380 T0	20-06-2002
			DE 60008360 D1	25-03-2004
			DE 60008360 T2	09-12-2004
			EP 1169301 A1	09-01-2002
			ES 2215672 T3	16-10-2004
			ES 2187390 A1	01-06-2003
			JP 2003522752 T	29-07-2003
			PL 350356 A1	02-12-2002
			RU 2225390 C2	10-03-2004
			WO 0158854 A1	16-08-2001
			WO 0158855 A1	16-08-2001
			ZA 200206497 A	14-08-2003
WO 0216310	A	28-02-2002	AU 8523001 A	04-03-2002
			BR 0114801 A	14-10-2003
			CA 2420279 A1	28-02-2002
			CN 1471508 A	28-01-2004
			EP 1401801 A1	31-03-2004
			EP 1491524 A2	29-12-2004
			JP 2004518617 T	24-06-2004
			WO 0216310 A1	28-02-2002
			US 2002099036 A1	25-07-2002
			US 2002099096 A1	25-07-2002
			US 2002173495 A1	21-11-2002
			US 2003022868 A1	30-01-2003
			US 2003162761 A1	28-08-2003
			US 2003232792 A1	18-12-2003
			US 2003225040 A1	04-12-2003
			US 2004014975 A1	22-01-2004
			US 2004260108 A1	23-12-2004
			US 2005038110 A1	17-02-2005
EP 0797980	A1	01-10-1997	DE 19612506 A1	02-10-1997
			BR 9701464 A	25-08-1998
			ES 2109207 T1	16-01-1998
			JP 9268170 A	14-10-1997
			US 5865856 A	02-02-1999
EP 0488756	A1	03-06-1992	JP 2701090 B2	21-01-1998
			JP 4202164 A	22-07-1992
			DE 69124780 D1	03-04-1997
			DE 69124780 T2	16-10-1997
			ES 2097798 T3	16-04-1997
			US 5238818 A	24-08-1993
US 4723986	A	09-02-1988	NONE	
JP 49020129	A2	22-02-1974	JP 49020129 A	22-02-1974
			CA 1012553 A1	21-06-1977
			CH 593236 A5	30-11-1977

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB2004/004464

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 49020129	A2	DE 2331900 A1	24-01-1974
		FR 2199533 A1	12-04-1974
		FR 2225423 A1	08-11-1974
		GB 1421600 A	21-01-1976
		IT 991637 B	30-08-1975
<hr/>			